

RESEARCH ARTICLE / ARAŞTIRMA MAKALESİ

A TIGHT BINDING MODEL FOR QUANTUM SPIN HALL EFFECT ON TRIANGULAR OPTICAL LATTICE

Ahad K. ARDABILI¹

¹Altınbaş University, School of Engineering and Natural Sciences,
Department of Basic Sciences, Istanbul.
ahad.ardabili@altinbas.edu.tr ORCID No: 0000-0002-0957-1833

Tekin DERELİ²

²Koç University, College of Sciences, Department of Physics, Istanbul.
tdereli@ku.edu.tr ORCID No: 0000-0002-6244-6054

Özgür E. MÜSTECAPLIOĞLU³

³Koç University, College of Sciences, Department of Physics, Istanbul.
omustecap@ku.edu.tr ORCID No: 0000-0002-9134-3951

Received Date/Geliş Tarihi: 07/11/2018. Accepted Date/Kabul Tarihi: 13/03/2019.

Abstract

We propose a tight binding model for the quantum spin Hall system on triangular optical lattice and we determined the edge state spectrum which contains gap traversing states as the hallmark of Z_2 topological insulator. The advantage of this system is the possibility of implementing it in the fermionic ultracold atomic system whose nearly free electron limit is proposed by B. Beri and N. R. Cooper, Phys. Rev. Lett. 107, 145301 (2011).

Keywords: Ultra cold atom, Topological insulator, Tight binding model, Quantum mechanics, Optical lattice.

ÜÇGEN OPTİK ÖRGÜDE KUANTUM SPİN HALL ETKİSİ İÇİN SIKI-BAĞ MODELİ**Öz**

Bu makede, üçgen örgü üzerinde kuantum spin Hall sistemi için bir sıkı-bağ modeli önermekte ve Z_2 topolojik yalıtkanının ayırt edici özelliği olan boşluk geçiş durumlarını da içeren kenar durum spektrumunu incelemekteyiz. Bu sistemin avantajı, serbest elektron sınırı B. Beri ve N. R. Cooper tarafından verilen (Phys. Rev. Lett. 107, 145301, 2011) fermiyonik aşırı soğuk atomik sistemlerde de incelenme imkanının olmasıdır.

Anahtar Kelimeler: Aşırı soğuk atomik sistemler, Topolojik yalıtkan, Bir sıkı-bağ modeli, Kuantum mekaniği, Optik örgü.

1. INTRODUCTION

Topological insulators (TIs) are insulating in the bulk but have metallic states on their boundaries [Hasan and Kane (2010), Zhang (2011)]. Robustness of these states against disorder and perturbations makes them promising for applications such as spintronics [Moore J. E. (2010)] and topological quantum computation [Nayak *et al.* (2008)]. Topological invariants of the bulk material are essential for the robust boundary modes. This urged consideration of topological insulators on different lattice geometries [Hu, Kargarian, and Fiete (2011), Weeks and Franz (2010), Guo and Franz (2009a), Guo and Franz (2009b), Kane, Fu and Mele (2007)].

It is widely acknowledged that the cold atomic systems are ideal systems to simulate solid-state phenomena in a controlled way. The two and three dimensional topological insulators with band gaps in the order of the recoil energy have recently been proposed in ultracold fermionic atomic gases [Beri and Cooper (2011)]. The proposal utilizes interactions which preserves time reversal symmetry (TRS), analogous to synthesized spin-orbit coupling [Lin, Jimenez-Garcia and Spielman (2011)], so that the insulators are classified by the so-called Z₂ topological invariant [Fu and Kane (2006)].

Even if the band gap in tight binding models are not as large as in nearly free electron limit, TIs in ultracold atomic systems have been studied vastly in tight binding regime [Juzelians, Ruseckas and Dalibard (2010), Goldman *et al.* (2010)]. The optical lattices are described by continuous potentials formed by the combinations of standing waves. It is convenient to treat them as deep potentials. Our aim in this article is to propose a tight binding model for the quantum spin Hall effect which can be realized in the ultracold atomic systems. The corresponding model in the nearly free electron limit is proposed by Beri and Cooper [Beri and Cooper (2011)] with this advantage that the band gap is large. We also determine the band structure of the edge state which exhibits the hallmark of TIs due to its robustness against all perturbations that preserve the TRS. In the Sec. II of this paper we propose the tight binding model for quantum spin Hall (QSH) system in the triangular optical lattice. In Sec. III we briefly review the proposal of Z₂ topological insulator in ultracold atomic gases [Beri and Cooper (2011)]. We conclude in Sec. IV.

2. TIGHT BINDING MODEL

2.1 Bulk band structure

The charge quantum Hall effect depends on the breaking of time-reversal symmetry and it has been shown that even in the absence of average non-zero external magnetic field the quantum Hall effect can be created [Haldane (1988)]. However, in the QSH effect one needs to preserve the time reversal invariance. Among the first models proposed for dissipationless QSH effect are the works by Bernevig and Zhang (2006) and by Kane and Mele (2005), where the authors used the spin-orbit coupling such that the two-different spin direction experiences the same magnetic field strength but with opposite sign. In other words, their system were two copies of a quantum Hall system for each spin where the total first Chern number adds up to zero and the system is time reversal invariant.

Physically our model corresponds to the same scenario. We propose a Hamiltonian for a fermion on triangular lattice Fig. 1 with a mirror symmetric spin orbit coupling as:

$$H = t \sum_{m,n} C_{m+1,n}^\dagger C_{m,n} + C_{m,n+1}^\dagger e^{i4\pi m\phi} \sigma_z C_{m,n} + C_{m+1,n-1}^\dagger e^{i2(m+1)\pi\phi} \sigma_z C_{m,n}, \quad (1)$$

where $\phi = p/q$ is flux per plaquette and we take $p = 1$ and $q = 4$ in this paper. $C_{m,n} = (c_{m,n\uparrow}, c_{m,n\downarrow})^T$ and $C_{m,n}^\dagger$ are annihilation and creation operators on site (m, n) respectively. We take the hopping parameter $t = 1$ throughout this paper. The first term is nearest neighbor hopping term on the triangular lattice with $\mathbf{a}_1 = (3/4, 1/4)\mathbf{a}$ and $\mathbf{a}_2 = (0, 1/2)\mathbf{a}$, where \mathbf{a} is the lattice constant (see Fig. 1). The second and third terms are mirror symmetric spin-orbit interaction. σ_z is the Pauli matrix. In the absence of spin this Hamiltonian implies that electron acquires $\phi = 1/4$ of flux quantum enclosing the elementary plaquette of the triangular lattice.




Figure 1. The light points and dashed lines show the lattice sites of the atoms and the hopping directions in the triangular lattice respectively, and the dark points and lines are the corresponding Brillouin zone sites and vectors respectively.

To calculate the band structure, we take the Fourier transform of the Hamiltonian Eq. (1). We use the momentum representation of fermionic operator

$$C_{\mathbf{k}} = \sum_{m,n} e^{i\mathbf{k}\cdot\mathbf{R}_{m,n}} C_{m,n}, \quad (2)$$

where $\mathbf{R}_{m,n} = m\mathbf{a}_1 + n\mathbf{a}_2$. We obtain the energy dispersion in triangular lattice by solving the determinant for the eigenvalues ϵ ,

$$\text{Det} \begin{bmatrix} -A - \epsilon & B + iC & 0 & 0 \\ B - iC & A - \epsilon & 0 & 0 \\ 0 & 0 & A - \epsilon & B - iC \\ 0 & 0 & B + iC & -A - \epsilon \end{bmatrix} = 0, \quad (3)$$

where A, B and C are defined to be:

$$\begin{aligned} A &= \cos(k_y a/2), \\ B &= \cos\frac{a}{4}(\sqrt{3}k_x + k_y), \\ C &= \cos\frac{a}{4}(\sqrt{3}k_x - k_y). \end{aligned} \quad (4)$$

To solve Eq. (3), we used a 2D grid for the k -space. Fig. 2a shows the band structure of the Eq. (3) for a cell with specific \mathbf{k} points as its corners taken to be $(\mathbf{k}_1 + \mathbf{k}_2)/2, k_1/2, 0, k_2/2$, as shown in the inset. These points are the TRS invariant points in the Brillouin zone. Since each of the two blocks of the Eq. (3) corresponds to two-fold spin degenerate bands, each band of the Fig. 2a is four-fold degenerate.




Figure 2. Left (a): The band structure of the Hamiltonian Eq. (1) along the path $(1,1), (1,0), (0,0), (0,1)$ where 1 and 0 are referring to the inset (m,n) in $\Gamma_{m,n} = (mk_1 + nk_2)$, which are the TRS invariant points in Brillouin zone. Right (b): The DOS for all the energy bands is shown where the horizontal axis is the energy.

We also calculated the density of states (DOS) for the Hamiltonian in Eq. (3). This quantity is defined by the expression:

$$\rho(E) = \frac{1}{A} \sum_{\mathbf{k}} \delta(E - E_n), \quad (5)$$

where A is the area of the system in reciprocal space and E_n is the energy of the bands. In Fig. 2b the DOS is depicted for all the energy bands.

2.2 Edge-state band structure

The characteristic of the Z2 topological insulator is the gapless edge states. They describe two spin currents at the edge, propagating in opposite direction. This property is because of the time-reversal symmetry and it prevents the gap opening due to any TR invariant perturbation as the result of the Kramer's theorem [Kane and Mele (2005)].

We follow the method in Ref. [Hatsugai (1993)] to find the energy dispersion of the edge states. The Hamiltonian Eq. (1), must be reduced to a one-dimensional problem. We take the y direction as the periodic part and we use the momentum representation as:

$$C_{m,n} = \frac{1}{\sqrt{L_y}} \sum_{k_y} e^{ik_y n} C_m(k_y), \quad (6)$$

where $\frac{k_y a}{2} = \frac{2\pi n_y}{L_y}$, $n_y = 1, \dots$, is the system size along y direction. By inserting the single particle state

$$|\Psi(k_y)\rangle = \sum_m |\Psi(k_y)_m\rangle C_m^\dagger(k_y)|0\rangle \quad (7)$$

into the Schrödinger equation $H|\Psi\rangle = E|\Psi\rangle$, the spin up part of the problem is reduced to the one-dimensional problem with parameter k_y as:

$$G^* \Psi_{m+1} - G \Psi_{m-1} - 2 \cos\left(\frac{k_y a}{2} - 4\pi\phi m\right) \Psi_m = E \Psi_m. \quad (8)$$

where $G = 1 + e^{-i(-k_y a/2 + \pi\phi(2m+1))}$. Including the spin down as well, this equation can be written as a generalized Harper equation [Harper (1955)] in transfer matrix form:

$$\begin{pmatrix} \Psi_{m+1\uparrow}(k_y) \\ \Psi_{m\uparrow}(k_y) \\ \Psi_{m+1\downarrow}(k_y) \\ \Psi_{m\downarrow}(k_y) \end{pmatrix} = M \begin{pmatrix} \psi_{m\uparrow}(k_y) \\ \psi_{m-1\uparrow}(k_y) \\ \psi_{m\downarrow}(k_y) \\ \psi_{m-1\downarrow}(k_y) \end{pmatrix}, \quad (9)$$

where M is the transfer matrix, which is given by:

$$M = \begin{pmatrix} \frac{F}{G} & \frac{G^*}{G} & & 0 \\ 1 & 0 & & \\ & & \frac{F'}{G'} & \frac{G'^*}{G'} \\ 0 & & 1 & 0 \end{pmatrix} \quad (10)$$

with

$$G' = 1 - e^{-i\left(-\frac{k_y a}{2} + \pi\phi(2m+1)\right)}, F = -\epsilon - 2 \cos\left(\frac{k_y a}{2} - 4\pi\phi m\right) \text{ and } F' = -\epsilon + 2 \cos(k_y a/2 - 4\pi\phi m).$$

Under the boundary condition that the wavefunction goes to zero at the boundaries of lattice we can solve this equation. The band structure along the path $k_y a = 0 - 2\pi$ is shown in the Fig. 3. We used 100 k-points along k_y direction. The shaded area is the bulk band and the gap traversing edge states as the signature of Z2 topological insulator are plotted as the solid line. Since the TRS is preserved no TR symmetric perturbation can open the gap at $k_y a = \pi$.




Figure 3. The spectrum of the edges of triangular lattice. The shaded area is lowest energy bands for the triangular lattice (for uniform spacing of $-4\pi/\sqrt{3} < k_x a < 4\pi/\sqrt{3}$). The solid lines show the spin polarized edge state band, traversing the gap. The edge states cross at

3. COLD ATOMIC SYSTEM

In this section, we review briefly the Z2 topological insulator model proposed by Berri and Cooper (2011). This model is studied in nearly free electron limit which has the advantage of large band gap.

The Hamiltonian which describes an atom with position \mathbf{r} and momentum \mathbf{p} and with N internal states is given by

$$H = \frac{\vec{p}^2}{2m} \mathbb{1}_4 + VM(\vec{r}) \quad (11)$$

where $VM(\vec{r})$ is a position dependent potential. To have a system with low spontaneous emission one can use ytterbium (Yb) which has long-lived excited state. The two internal states, ground state ($1S_0 = g$) and long-lived excited state ($3p_0 = e$) of Yb have spin degree of freedom which leads to four states. Another interesting aspect of Yb is the existence of a state dependent scalar potential for λ_{magic} with opposite sign $\pm V_{\text{am}}(\mathbf{r})$. Therefore, we can write the potential part of Hamiltonian when we have external electric field as $\mathbf{E} = \epsilon e^{-i\omega t} + \epsilon^* e^{i\omega t}$ with complex amplitude ϵ and frequency ω . All four e-g transitions have the same frequency $\omega_0 = (E_e - E_g)/\hbar$. Using rotating wave approximation [Cohen-Tannoudji, Dupont-Roc, and Grynberg (1992)] we have the optical potential as following:

$$VM(\vec{r}) = \begin{pmatrix} (\frac{\hbar}{2}\Delta + V_{\text{am}})\mathbb{1} & -i\vec{\sigma} \cdot \vec{\epsilon} d_r \\ i\vec{\sigma} \cdot \vec{\epsilon} d_r & -(\frac{\hbar}{2}\Delta + V_{\text{am}}) \end{pmatrix} \quad (12)$$

where $\Delta = \omega - \omega_0$ is the atom-field detuning and d_r is the dipole moment. One can write the Hamiltonian in terms of Dirac matrices [Murakami, Nagaosa, and Zhang (2003)]:

$$\Gamma^{1,2,3} = -i\sigma^y \otimes \sigma^i, \Gamma^4 = \sigma^x \otimes I \text{ and } \Gamma^5 = \sigma^z \otimes I, \quad (13)$$

which gives:

$$H = \frac{\hat{P}^2}{2m} \mathbf{1} + \Gamma^i d_r \epsilon_i + \Gamma^5 \left(\frac{\hbar}{2} \Delta + V_{am} \right), \quad (14)$$

For the two-dimensional system one can make following choice for the potential matrix Eq. (12):

$$\begin{aligned} d_r \epsilon &= [V\delta, V\cos(\vec{r} \cdot \vec{k}_1), V\cos(\vec{r} \cdot \vec{k}_2)], \\ \frac{\hbar}{2} \Delta + V_{am}(\vec{r}) &= V\cos(\vec{r} \cdot (\vec{k}_1 + \vec{k}_2)) \end{aligned} \quad (15)$$

with $\mathbf{k}_1 = k(1,0,0)$ and $\mathbf{k}_2 = k(\cos(\theta), \sin(\theta), 0)$. The optical potential in Eq. (17) is formed from three standing waves which are linear polarized light at the coupling frequency ω . Two of these waves have equal amplitude in the 2D plane (\mathbf{k}_1 for y polarization and \mathbf{k}_2 for z polarization) the x-polarized wave vector is normal to the 2D plane with an amplitude smaller by a factor of δ . Since the $\cong \omega_0$, we have $k \cong 2\pi/\lambda_0$ with $\lambda_0 = 578 \text{ nm}$ the wavelength of the e-g transition. The spatial dependence of V_{am} is set by a standing wave at the antimagic wavelength λ_{am} [Gerber and Dalibard (2010)], which creates a state-dependent potential with $|\mathbf{k}_1 + \mathbf{k}_2| = 4\pi/\lambda_{am}$ that leads $\theta = 2 \arccos(\pm\lambda_0/\lambda_{am})$. For simplicity, in all following discussions one can fix $\theta = 2\pi/3$ and define $a \equiv 4\pi/(\sqrt{3}k)$. Therefore the optical coupling M^* has the symmetry of a triangular lattice. In Fig. 4a we show the few lowest energy bands for $\delta = 0$. The bands were calculated by numerical diagonalization in the plane wave basis (49 plane waves are used). All bands are fourfold degenerate like tight binding regime. The relation of this system to the tight binding model given in previous section becomes clearer as one applies the unitary transformation $\hat{U} = (1 - i\hat{\Sigma}_3\hat{\sigma}_2)/\sqrt{2}$ to the coupling \hat{M} in Eq. (12):

$$\hat{M}' = \hat{U}^\dagger \hat{M} \hat{U} = c_1 \hat{\Sigma}_1 + c_2 \hat{\Sigma}_2 \sigma_3 + c_{12} \hat{\Sigma}_3 \quad (16)$$

here $\Sigma_i = \sigma_i \otimes 1_{2 \times 2}$, $c_i \equiv \cos(\mathbf{r} \cdot \mathbf{k}_i)$ and $c_{12} = \cos(\mathbf{r} \cdot (\mathbf{k}_1 + \mathbf{k}_2))$. This matrix is 2×2 block diagonal matrix for each eigenvalue of σ_3 (since the kinetic part is diagonal this is the case for the Hamiltonian as well) thus the four-level system decouples into two two level system each of which experiences an effective magnetic field due to the optical dressed state of the $c_1 \hat{\Sigma}_1 \pm c_2 \hat{\Sigma}_2 + c_{12} \hat{\Sigma}_3$ [Cooper (2011)]. This means that opposite spin direction undergoes an effective magnetic field of the

same strength but with opposite signs. Beside the lowest band energy of these systems have ± 1 Chern number which they cancel out each other because of the time-reversal symmetry. These are the required criteria for the quantum spin Hall effect.


To make the connection to the previous section we consider the adiabatic limit $V \gg \hbar k^2/2M$ [Jaksch and P. Zoller (2005)] when the potential part of the Hamiltonian Eq. (11) plays the dominant role. To find the minima of the adiabatic energy which gives the lattice sites in tight binding regime [Jaksch *et al.* (1998)] we diagonalized the potential in Eq. (16) analytically and obtained:

$$V_{ad} = \pm \sqrt{c_1^2 + c_2^2 + c_{12}^2}, \tag{17}$$

as plotted in Fig. 5.

Now if we ignore the spin here, the effective magnetic field strength experienced by the neutral atom following the adiabatic path is equivalent to the 1/4 of a flux quantum that a charged particle acquires enclosing the elementary plaquette of the triangular lattice in tight binding limit [Ashcroft and Mermin (1976)]. This is equivalent to the Hamiltonian Eq. (1) proposed in this paper without spin. Therefore, tight binding limit of the ultra-cold atomic system of Ref. [Beri and Cooper (2011)] is given in Eq. (1) and the hopping parameter t is related to the potential scale of optical coupling V based on the formalism in [Jaksch, *et al.* (1998)]. The difference is in the (0, 0) point in k -space where the upper band of cold atom limit has a sharper peak Fig. 4a than the tight binding upper band Fig. 2a. This can be understood as the characteristic behavior of the energy levels of free electron which are just a parabola in k (momentum), by getting distorted due to a periodic potential [Ashcroft and Mermin (1976)].

As the potential becomes stronger the energy dispersion resembles the tight binding regime Fig. 4b. The DOS for the nearly free electron limit is also depicted in Fig. 4c which shows that states are distributed around the two energy bands across the gap asymmetrically due to the asymmetrically located van Hove singularities of the upper and lower bands in contrast to that of tight binding case in Fig. 2b. Finally, we note that realization of QSH models in Eq. (1) and Eq. (11) does not require any spin flipping interactions and thus does not need any additional cooling mechanism [Kennedy *et al.* (2013)].






Figure 4. Top Left(a): Lowest energy bands of the cold atom system in nearly free electron limit as the result of the solution (a). Lowest energy bands of the cold atom system in nearly free electron limit as the result of the solution of Eq. (14). The energy is plotted relative to recoil energy E_R . Here $V = 0.5E_R$ and $\delta = 0$ are shown here. The k-points are labeled as $\Gamma_{mn} = (mk_1 + nk_2)/2$ and each band is four-fold degenerate.

Bottom left (b): Energy bands for the potential $V = 3.5E_R$ which resembles the band structure of tight binding regime Fig. 2a.

Right (c): DOS of the energy bands in Fig. 2a which are depicted in the band structure is shown. Energy is expressed in the unit of E_R .




Figure 5. The dark circles show the local minima of the adiabatic energy which forms a triangular lattice in the tight binding limit.

4. CONCLUSION

Summarizing, we considered the quantum spin Hall effect on the triangular lattice in the tight binding limit and we proposed that this can be realized in the ultra-cold atomic system. We studied the edge state band structure which reveals the Z2. The nearly free electron limit of the system we proposed here is introduced as Z2 topological insulator in Ref. [Beri and Cooper (2011)].

Acknowledgments

Authors acknowledge useful discussions with O. Oktel and I. Adagideli.

Conflict of Interests/Çıkar Çatışması

Authors declare no conflict of interests/Yazarlar çıkar çatışması olmadığını belirtmişlerdir.

5. REFERENCES

- Ashcroft N. W. and N. D. Mermin.** 1976. "Solid State Physics", Thomson Learning, Toronto.
- Beri B. and N. R. Cooper.** 2011. "Z₂ Topological Insulators in Ultracold Atomic Gases", Phys. Rev. Lett., 107, 145301.
- Bernevig B. A. and S.-C. Zhang.** 2006. "Quantum Spin Hall Effect", Phys. Rev. Lett., 96, 106802.
- Cohen-Tannoudji C., J. Dupont-Roc, and G. Grynberg,** 1992. "Atom-Photon Interactions", Wiley, New York.
- Cooper N.** 2011. "Optical Flux Lattices for Ultracold Atomic Gases", Phys. Rev. Lett., 106, 175301 2.
- Dalibard J., F. Gerbier, G. Juzeliūnas, and P. Öhberg.** 2011. "Artificial gauge potentials for neutral atoms" Rev.Mod., Phys. 83, 1523.
- Fu L. and C. L. Kane.** 2006. "Time reversal polarization and a Z₂ adiabatic spin pump", Phys. Rev. B, 74, 195312.
- Gerbier F. and J. Dalibard.** 2010. "Gauge fields for ultracold atoms in optical superlattices", New Jour. Phys., 12, 033007.
- Goldman N, I. Satija, P. Nikolic, A. Bermudez, M. A. Martin-Delgado, M. Lewenstein, and I. B. Spielman.** 2010. "Realistic Time-Reversal Invariant Topological Insulators with Neutral Atoms", Phys. Rev. Lett., 105, 255302.
- Guo H.-M. and M. Franz.** 2009. "Topological insulator on the kagome lattice", Phys. Rev. B, 80, 113102.
- Guo H.-M and M. Franz.** 2009. "Three-Dimensional Topological Insulators on the Pyrochlore Lattice", Phys. Rev. Lett., 103, 20680.
- Haldane F. D. M.** 1988. "Model for a Quantum Hall Effect without Landau Levels: Condensed-Matter Realization of the "Parity Anomaly", Phys. Rev. Lett., 61, 2015.
- Harper P. G.** 1955. "The General Motion of Conduction Electrons in a Uniform Magnetic Field, with Application to the Diamagnetism of Metals", Proc. Phys. Soc. A, 68, 874.
- Hasan M. Z. and C. L. Kane.** 2010. "Colloquium: Topological insulators", Rev. Mod. Phys., 82, 3045.

- Hatsugai Y.** 1993. "Edge states in the integer quantum Hall effect and the Riemann surface of the Bloch function", Phys. Rev. B, 48, 11851.
- Hu X., M. Kargarian, and G. A. Fiete.** 2011. "Topological Insulators and fractional quantum Hall effect on the ruby lattice", Phys. Rev. B, 84, 155116.
- Jaksch D. and P. Zoller.** 2005. "The cold atom Hubbard toolbox", Annals of Physics, 315, 52.
- Jaksch D., C. Bruder, J. I. Cirac, C. W. Gardiner, and P. Zoller.** 1998. "Cold Bosonic Atoms in Optical Lattices", Phys. Rev. Lett., 81, 3108.
- Juzeliūnas G., J. Ruseckas, and J. Dalibard.** 2010. "Generalized Rashba-Dresselhaus spin-orbit coupling for cold atoms", Phys. Rev. A, 81, 053403.
- Kane C. L., L. Fu and E. J. Mele.** 2007. "Topological Insulators in Three Dimensions", Phys. Rev. Lett., 98, 106803.
- Kane C. and E. Mele.** 2005. " Z_2 Topological Order and The Quantum Spin Hall Effect", Phys. Rev. Lett., 95, 146802.
- Kane C. and E. Mele.** 2005. "Quantum Spin Hall Effect in Graphene", Phys. Rev. Lett., 95, 226801.
- Kennedy C. J., G. A. Siviloglou, H. Miyake, W. C. Burton, and W. Ketterle.** 2013. "Spin-Orbit Coupling and Quantum Spin Hall Effect for Neutral Atoms without Spin Flips", Phys. Rev. Lett., 111, 225301.
- Lin Y.-J., K. Jimenez-Garcia, and I. B. Spielman.** 2011. "Spin-orbit-coupled Bose-Einstein condensates", Nature, 471, 83.
- Moore J. E.** 2010. "The birth of topological insulators", Nature, 464, 194.
- Murakami S., N. Nagaosa, and S. Zhang.** 2003. "Dissipationless Quantum Spin Current at Room Temperature", Science, 301, 1348.
- Nayak C., S. H. Simon, A. Stern, M. Freedman, and S. D. Sarma.** 2008. "Non-Abelian anyons and topological quantum computation", Rev. Mod. Phys., 80, 1083.
- Weeks C. and M. Franz.** 2010. "Topological insulators on the Lieb and perovskite lattices", Phys. Rev. B, 82, 085310, 2010.
- Zhang S. C.** 2011. "Topological insulators and superconductors", Rev. Mod. Phys., 83, 1057.

