

MODELING DEPENDENT FINANCIAL ASSETS BY DYNAMIC COPULA AND PORTFOLIO OPTIMIZATION BASED ON CVAR

SIBEL AÇIK KEMALOĞLU AND EMEL KIZILOK KARA

ABSTRACT. This paper is concerned with the statistical modeling of the dependence structure of multivariate financial data using copula. Since financial data is greatly affected by the economic factors, it often varies according to the time. Therefore, dynamic copula model is used that takes into account the time-varying. In addition, portfolio optimization based on Mean-CVaR model is applied with Monte Carlo simulation. As an application, a portfolio with four different Indexes is constructed from the Turkish financial markets. The marginal distributions of assets in the portfolio are estimated and parameter estimates are given for the different copula models. The portfolio optimization based on CVaR is made for the portfolio created from the specified copula model.

1. INTRODUCTION

Modern Portfolio Theory is an approach which had emerged after the 1950s and has enabled the creation of portfolios, taking into account the relationships between assets. Markowitz [1] presented the mean variance model and laid the foundations of modern portfolio theory. Portfolio risk is measured by variance in the mean variance model. When the return data is not distributed normally, it is more appropriate to use other risk measurements instead of measuring risk by variance. Value at Risk (VaR) and Conditional Value at Risk (CVaR) are the most known risk measurements in the field of finance. In the first studies carried out in finance, VaR was used as risk measurement. However, VaR could not ensure the attributes of being subadditive and convex, which are required in a risk measurement. It is of importance to ensure these attributes, particularly in a portfolio optimization. CVaR has been recommended as opposed to VaR, since it can ensure these attributes [2]. In this study, risk measurement was adopted as CVaR, and the mean-CVaR model was used for portfolio optimization.

In order to estimate the portfolio CVaR in a more accurate way, the nonlinear dependency between the tails of return on assets should be revealed. In this study,

Received by the editors: October 21, 2014 Accepted: January 29, 2015.

Key words and phrases. Dynamic copula, CVaR, portfolio optimization.

the copula model that was first mentioned in Sklar [3] was used in order to model the dependency structure between the return series. However, since the financial data exploited varied with respect to time, the dynamic copula model was applied to the data instead of a known copula model [4].

In order to conduct a portfolio optimization based on CVaR, the copula parameter should primarily be estimated. In the literature, there are different methods for the copula parameter estimation. The Inference Function for Margins (IFM) method, based on the marginal distribution, Kendall's τ based on sequence independent of marginal functions and Maximum Pseudo Likelihood (MPL) method, are the most known estimation methods. In this study, Two-Stage IFM method was used. In this method, the parameter of marginal distribution is estimated in the first phase, and copula parameter is estimated in the second phase [5]. For the test of marginal and copula goodness of fit, AIC and BIC criteria were applied. Finally, portfolio optimization based on the CVaR risk measurement was made for the data generated from this model with the Monte Carlo simulation method, and the weights in the portfolio were determined.

Comprehensive information regarding copula is available in the study of Joe [6] and Nelsen [7].

In the field of finance, copula was first mentioned by Embrechts et al. [8]. The first applications of this subject were given in Cherubini [5]. Jondeau and Rockinger [9] used the Copula-GARCH model in order to determine the dependence structure among stocks. Other studies that included this model have been presented in Wei and Zhang [10], Ozun and Cifter [11], and Huang et al. [12]. Optimization studies based on CVaR were recommended in Uryasev and Rockafellar [2] for the first time. The application of the copula-GARCH model for portfolio risk analysis was given in Wu and Chen [13], and Wang et al. [14]. He and Li [4] have conducted studies regarding the dynamic copula model based on CVaR and copula.

In the second part of the study, the dynamic copula model is introduced, and methods to find the estimations for parameters are given. In the third part, the definition of CVaR risk measurement is presented and portfolio optimization based on CVaR is formed. In the fourth part, for a sample of Turkish financial data is tested and the results are presented. Finally, in the fifth part, the results are discussed and further studies that could be conducted in the future are mentioned.

2. DYNAMIC COPULA MODEL AND PARAMETER ESTIMATION

Copula functions are frequently used in modeling the dependency structure among the risk variables in the field of finance and actuary. Sklar [3] states that there is only one expression of an n -dimensional $C(\cdot, \dots, \cdot)$ copula for any continuous (X_1, \dots, X_n) random vector:

$$F(x_1, \dots, x_n) = C(F_1(x_1), \dots, F_N(x_n)) \quad (2.1)$$

where $F_1(\cdot), \dots, F_N(\cdot)$ and $F(\cdot, \dots, \cdot)$ respectively show the marginal and joint distribution functions of x_1, x_2, \dots, x_n random variables.

In this study, static copula models (Gaussian, Student-t, Clayton, and Symmetrized Joe Clayton-SJC) and corresponding dynamic copula models (GDCC, tDCC, tvC, tvSJC) are used. The static copulas exploited are defined as follows [5, 7, 15]

Table 1. Bivariate copula functions and parameter space of the considered copulas

Copula	$C(u_1, u_2)$	Parameter Space
Gaussian	$\Phi_\rho(\phi^{-1}(u_1), \phi^{-1}(u_2)) = \int_{-\infty}^{\phi^{-1}(u_1)} \int_{-\infty}^{\phi^{-1}(u_2)} \frac{1}{2\pi\sqrt{1-\rho^2}} \exp\left(-\frac{s^2-2\rho st+t^2}{2(1-\rho^2)}\right) ds dt$	$\rho \in (-1, 1)$
Student's t	$t_{d,\rho}(t_d^{-1}(u_1), t_d^{-1}(u_2)) = \int_{-\infty}^{t_d^{-1}(u_1)} \int_{-\infty}^{t_d^{-1}(u_2)} \frac{1}{2\pi\sqrt{1-\rho^2}} \left(1 + \frac{s^2-2\rho st+t^2}{d(1-\rho^2)}\right) ds dt$	$\rho \in (-1, 1)$ $d \in (0, \infty)$
Clayton	$[u_1^{-w} + u_2^{-w} - 1]^{-\frac{1}{w}}$	$w \in (0, \infty)$
SJC	$\frac{1}{2} (C_{JC}(u_1, u_2 \tau^U, \tau^L) + C_{JC}(1 - u_1, 1 - u_2 \tau^L, \tau^U) + u_1 + u_2 - 1)$	Upper tail: $\tau^U \in (0, 1)$ Lower tail: $\tau^L \in (0, 1)$

Also note that ϕ^{-1} and t_d^{-1} are the inverse of the Normal and t-student c.d.f., where the parameters ρ and d are the coefficient of linear correlation and the degrees of freedom, respectively. Joe-Clayton copula is defined as $C_{JC}(u_1, u_2 | \tau^U, \tau^L) = 1 - \left(1 - \{[1 - (1 - u_1)^\kappa]^{-\gamma} + [1 - (1 - u_2)^\kappa]^{-\gamma} - 1\}^{-1/\gamma}\right)^{1/\kappa}$, where $\kappa = \frac{1}{\log_2(2-\tau^U)}$ and $\gamma = \frac{1}{\log_2(\tau^L)}$

The used dynamic copula are created by applying the static copulas which are defined in Table 1. According to the Sklar [3] theorem, the dynamic copula model is expressed as below [11].

$$F(X_{1t}, \dots, X_{nt} | \xi_t) = C_t(F_{1t}(X_{1t} | \xi_t), F_{2t}(X_{2t} | \xi_t), \dots, F_{nt}(X_{nt} | \xi_t)) \quad (2.2)$$

where $\xi_t = \sigma\{X_{1t-1}, X_{2t-2}, \dots, X_{nt-t}, \dots\}$ for $t = 1, 2, \dots, T$ represents the historical data until t time.

The first step to create the joint distribution of portfolio assets with the dynamic copula method is generating the marginal distribution of each asset. In the literature, since the financial data varies according to the time, it is stated that marginal distributions are in compliance with the GARCH model, which was first run by Bollerslev [16]. The GJR model was acquired by adding the dummy variable to the GARCH model.

$GARCH(1,1) - n$ and $GARCH(1,1) - t$ models can be expressed as below:

$$\begin{aligned} x_t &= \mu + a_t \\ a_t &= \sigma_t \varepsilon_t \\ \sigma_t^2 &= \alpha_0 + \alpha_1 a_{t-1}^2 + \beta \sigma_{t-1}^2 \\ \varepsilon_t &\sim N(0,1) \text{ or } \varepsilon_t \sim t_d \end{aligned} \quad (2.3)$$

Here, provided that μ indicates conditional mean of return series, and σ_{t-1}^2 indicates conditional variance, it is defined as below. Moreover, Ω_{t-1} indicates the information set and d indicates the degree of freedom.

$$\begin{aligned} \mu &= E(x_t) = E(E(x_t|\Omega_{t-1})) = E(\mu_t) = \mu \\ \sigma_t^2 &= Var(x_t|\Omega_{t-1}) = Var(a_t|\Omega_{t-1}) \\ \alpha_0 &> 0, \alpha_1 \geq 0, \beta \geq 0 \text{ ve } \alpha_1 + \beta < 1 \end{aligned}$$

GJR-n and GJR- t models are defined as:

$$\begin{aligned} x_t &= \mu + a_t \\ a_t &= \sigma_t \varepsilon_t \\ \sigma_t^2 &= \alpha_0 + \alpha_1 a_{t-1}^2 + \beta \sigma_{t-1}^2 + \gamma s_{t-1} a_{t-1}^2 \\ \varepsilon_t &\sim N(0,1) \text{ or } \varepsilon_t \sim t_d \end{aligned} \quad (2.4)$$

where, $s_{t-1} = \begin{cases} 1, & a_{t-1} < 0 \\ 0, & a_{t-1} \geq 0 \end{cases}$ is a dummy variable which is equal to 0 as opposed to 1 while ε_t is negative. Moreover, parameter intervals are given as $\alpha_0 > 0, \alpha_1 \geq 0, \beta \geq 0, \beta + \gamma \geq 0$ and $\alpha_1 + \beta + \frac{1}{2}\gamma < 1$.

The parameters of the GARCH and GJR models are estimated with MLE method. The joint density function can be expressed as

$$f(a_1, \dots, a_t) = f(a_t|\Omega_{t-1}) f(a_{t-1}|\Omega_{t-2}) \dots f(a_1|\Omega_0) f(a_0)$$

where $\Omega_{t-1} = \{a_0, a_1, \dots, a_{t-1}\}$. Given data a_1, \dots, a_t , the log-likelihood function is given as:

$$LL = \sum_{k=0}^{n-1} f(a_{n-k}|\Omega_{n-k-1})$$

Here, the conditional marginal distribution of X_{t+1} for the $GARCH(1,1)$ model is defined as follows:

$$\begin{aligned} P(X_{t+1} \leq x|\Omega_t) &= P(a_{t+1} \leq (x - \mu) |\Omega_t) \\ &= P\left(\varepsilon_{t+1} \leq \frac{(x - \mu)}{\sqrt{\alpha_0 + \alpha_1 a_t^2 + \beta \sigma_t^2}} |\Omega_t\right) \\ &\begin{cases} N\left(\frac{(x - \mu)}{\sqrt{\alpha_0 + \alpha_1 a_t^2 + \beta \sigma_t^2}} |\Omega_t\right), & \varepsilon \sim N(0,1) \\ t_d\left(\frac{(x - \mu)}{\sqrt{\alpha_0 + \alpha_1 a_t^2 + \beta \sigma_t^2}} |\Omega_t\right), & \varepsilon \sim t_d \end{cases} \end{aligned}$$

For the $GJR(1, 1)$ model:

$$\begin{aligned} P(X_{t+1} \leq x | \Omega_t) &= P(a_{t+1} \leq (x - \mu) | \Omega_t) \\ &= P\left(\varepsilon_{t+1} \leq \frac{(x - \mu)}{\sqrt{\alpha_0 + \alpha_1 a_t^2 + \beta \sigma_t^2 + \gamma s_t a_t^2}} | \Omega_t\right) \\ &\quad \begin{cases} N\left(\frac{(x - \mu)}{\sqrt{\alpha_0 + \alpha_1 a_t^2 + \beta \sigma_t^2 + \gamma s_t a_t^2}} | \Omega_t\right), & \varepsilon \sim N(0, 1) \\ t_d\left(\frac{(x - \mu)}{\sqrt{\alpha_0 + \alpha_1 a_t^2 + \beta \sigma_t^2 + \gamma s_t a_t^2}} | \Omega_t\right), & \varepsilon \sim t_d \end{cases} \end{aligned}$$

The copula parameter should also be estimated in order to conduct portfolio optimization based on CVaR. Therefore, the IFM method that is composed of two phases based on the method of Maximum Likelihood Estimation (MLE) was used. The IFM estimates the marginal distribution parameters separately from the copula parameters. The estimation procedure of the IFM method consists of two steps [5].

In the first step, the parameters of the marginals are estimated:

$$\hat{\theta}_1 = \arg \max_{\theta_1} \sum_{t=1}^T \sum_{j=1}^n \ln f_j(x_{jt}; \theta_1). \quad (2.5)$$

In the second step, the parameter of the copula model is estimated, given θ_1 :

$$\hat{\theta}_2 = \arg \max_{\theta_2} \sum_{t=1}^T \ln c\left(F_1(x_{1t}), F_2(x_{2t}), \dots, F_n(x_{nt}); \theta_2, \hat{\theta}_1\right). \quad (2.6)$$

The IFM estimator is defined as (Cherubini et.al., [5])

$$\hat{\theta}_{IFM} = \left(\hat{\theta}_1, \hat{\theta}_2\right)' \quad (2.7)$$

To make the goodness of fit tests for marginal and copula, AIC (Akaike Information Criterion) [17] and BIC (Bayesian Information Criterion) [18, 19] criteria were used. AIC and BIC values are,

$$AIC = -2 * LL + 2k \quad (2.8)$$

$$BIC = -2 * LL + \ln(n) * k \quad (2.9)$$

where LL is the log-likelihood at its maximum point of the model estimated and k is the number of copula parameters in the model. According to these criteria, the best choice is the model with minimum AIC or BIC value.

3. PORTFOLIO OPTIMIZATION BASED ON CVAR

CVaR measurement was introduced to the literature by the development of VaR measurement by Rockafellar and Uryasev [2]. Let $x = (x_1, x_2, \dots, x_n)^T$ vector be a

portfolio with n assets, and let $y = (y_1, y_2, \dots, y_m)^T$ be m type loss factor with $p(y)$ density distribution. The loss factor of portfolio $f(x, y)$ is given as follows:

$$\Psi(x, \alpha) = \int_{f(x, y) \leq \alpha} p(y) dy, \quad \alpha \in R$$

For the given $\beta \in (0, 1)$ confidence level

$$\begin{aligned} \alpha_\beta(x) &= \min \{ \alpha \in R : \Psi(x, \alpha) \geq \beta \} \\ \Phi_\beta(x) &= E [f(x, y) | f(x, y) \geq \alpha_\beta(x)] \\ &= \frac{1}{1 - \beta} \int_{f(x, y) \geq \alpha_\beta(x)} f(x, y) p(y) dy \end{aligned}$$

$\alpha_\beta(x)$ and $\Phi_\beta(x)$ show *VaR* and *CVaR* at β confidence level, respectively. Rockafellar and Uryasev [2] converted the minimization problem of *CVaR* to that of $F_\beta(x, \alpha)$. Here, $F_\beta(x, \alpha)$ is a constantly differentiable function, which is composed of convex composition of *VaR* and *CVaR*.

$$F_\beta(x, \alpha) = \alpha + \frac{1}{1 - \beta} \int_{y \in R^m} [f(x, y) - \alpha]^+ p(y) dy \quad (3.1)$$

Since the analytical expression of $p(y)$ is difficult, or it is not known generally in practice, y 's are derived through the Monte Carlo simulation method. Accordingly, the (3.1) equation can be written as follows:

$$\tilde{F}_\beta(x, \alpha) = \alpha + \frac{1}{m(1 - \beta)} \sum_{j=1}^m [-x^T y_j - \alpha]^+ \quad (3.2)$$

Here, provided that m indicates the number of simulations, $x = (x_1, x_2, \dots, x_n)$ indicates the weights of assets in the portfolio, and $y_j = (y_{j1}, y_{j2}, \dots, y_{jn})$ indicates the derived returns, the portfolio selection model based on *CVaR* optimization is given as below:

$$\begin{aligned} \min \tilde{F}_\beta(x, \alpha) &= \alpha + \frac{1}{m(1 - \beta)} \sum_{j=1}^m z_j \\ z_j &= [-x^T y_j - \alpha]^+ \\ &\left\{ \begin{array}{l} x^T y_j + \alpha + z_j \geq 0 \\ z_j \geq 0 \\ \frac{1}{q} x^T \sum_{j=1}^m y_j \geq \rho \\ \sum_{i=1}^m x_i = 1, \quad x \geq 0 \end{array} \right\} \quad (3.3) \end{aligned}$$

Here, ρ is the return expected by investor. *CVaR* optimization problem can be solved as a linear programming problem. Returns generated from the dynamic copula model are obtained from the Monte Carlo simulation.

4. NUMERICAL EXAMPLE

In this study, in order to demonstrate the performance of the copula-GARCH model, 1649 daily stock final quotations between January 4, 2007 and August 1, 2013, as well as USD and EURO currency data, were used. Stock final quotations were obtained from the website of BIST [20], and USD and EURO currency data were retrieved from the website of the Turkish Central Bank [21]. Daily market returns of BIST30, BIST100, USD and EURO are shown in Figure 1.

When r_t indicates the returns, it is given as:

$$r_t = 100 \times \ln \left(\frac{P_t}{P_{t-1}} \right)$$

Here P_t indicates the index value at t time. The existence of volatility clustering is seen in Figure 1. Therefore, the results are given in Table 2, after quadratic returns that are named as ARCH effects are tested with the Engle test (LM(4), LM(6), LM(8), and LM(10)) to reveal whether there is a relation to the series.




Figure 1. Daily returns of BIST30, BIST100, USD and EURO




Figure 2. Autocorrelations of BIST30, BIST100, USD and EURO indexes (a) and partial autocorrelation measurements (b)

Table 2. Descriptive statistics and tests

Statistics	BIST30	BIST100	USD	EURO				
Sample Number	1646	1646	1646	1646				
Mean	0.000462	0.000479	0.000191	0.000195				
Standart deviation	0.0200	0.0185	0.0090	0.0081				
Skewness	-0.1174	-0.2545	-0.5112	0.2224				
Kurtosis	3.7106	4.1608	24.03122	7.4139				
Tests	Q-stat	p-value	Q-stat	p-value	Q-stat	p-value	Q-stat	p-value
Jarque-Bera	940.51	0.0010	1195.9	0.0010	39427	0.0010	3756.8	0.0010
LM(4)	118.5178	0.0000	124.0513	0.0000	227.9964	0.0000	191.0501	0.0000
LM(6)	129.5706	0.0000	132.7788	0.0000	228.3458	0.0000	199.3508	0.0000
LM(8)	160.8098	0.0000	152.2154	0.0000	286.3049	0.0000	258.9005	0.0000
LM(10)	166.1320	0.0000	155.6451	0.0000	287.7057	0.0000	268.2120	0.0000

Table 2 consists of summary statistics of financial returns and statistical test information in relation to ARCH effects. Thus, it is revealed that BIST30, BIST100, and USD have negative skewness (-0.1174, -0.2545 and -0.5112, respectively), the EURO has a positive skewness (0.2224), and marginal are not distributed normally according to JB statistics ($p\text{ value} < 0.05$). It should be looked to the Engle test results to determine whether there is correlation between the data related to each financial return. According to the Engle test results, LM statistics indicate that there are ARCH effects ($p\text{ value} < 0.05$). Furthermore, the correlations for raw returns in Figure 2 were evaluated with autocorrelation functions (ACF) and partial autocorrelation functions (PACF). Here, it was observed that all autocorrelations are not zero after zero lag. Since financial returns are considered to be time series data, their marginal distributions were regarded as GARCH and GJR models to conduct copula estimation. Another result that supports the selection of these

models is that kurtosis coefficients that were computed for each of the financial returns were found to be greater than 3. As a consequence, empiric observations in regard to returns have greater leptokurtosis than the normal distribution. Goodness of fit for the distribution with regard to the residual series of each of the financial returns (BIST30, BIST100, USD, and EURO) were made for the GARCH and GJR models by selecting Normal and Student-t, respectively.

Table 3 and Table 4 respectively show maximum likelihood results, estimated parameter values, AIC (Akaike Information Criterion) and BIC (Bayesian Information Criterion) values for GARCH and GJR models selection.

In order to determine the model that best explains financial returns, minimum AIC and BIC values should be sought in the tables. Table 3 and Table 4 demonstrate that the GJR-t (AIC:-8612.8174,-8895.0195) model can be used for BIST 30 and BIST 100, and the GARCH-t (AIC: -11681.6623, -11889.9500) model can be used for USD and EURO.

An appropriate copula model was selected by using the estimated parameter values for the identified marginal distributions. Four copula functions were created for the application. The copula selection model was examined in two situations: static (Gaussian, Student-t, Clayton, SJC) and dynamic (GDCC, tDCC, tvC, tvSJC). Table 5 demonstrates the parameter estimation results for the IFM method in regard to these situations, and AIC and BIC values. From the table, it is understood that in both situations, the Student-t copula is the best copula model in order to explain the dependency structure of financial returns with four variables. When examined separately for static and dynamic situations, the dependency structure of GJR marginal distributions with the minimum AIC (-8831.6529) in a dynamic situation was modeled ideally with tDCC copula.

It was identified that financial data marginals that were used fit GJR-t, and joint dependency structure fit the tDCC dynamic copula model. By using the parameter values for selected model, the Monte Carlo simulation was performed 10,000 times, and portfolio optimization was achieved based on CVaR risk measurement. VaR and CVaR risk measurement values for different confidence levels, and results including the weights for each financial asset are given in Table 6. Accordingly, mean loss above a certain value, that is, CVAR value would reach the minimum when an investor allocates 35% of his/her assets in BIST30, 15% in BIST100, 30% in USD, and 20% in EURO within a 99% confidence level.

Table 3. Parameter estimates of GARCH-n and GARCH-t models and statistic tests

GARCH-n								
	BIST30		BIST100		USD		EURO	
Parameter	Value	Std	Value	Std	Value	Std	Value	Std
μ	0.0015	0.0000	0.0016	0.0000	0.0001	0.0000	0.0000	0.0000
α_0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
α_1	0.1198	0.0320	0.1325	0.0380	0.0994	0.0150	0.1028	0.0400
β	0.8548	0.0390	0.8452	0.0410	0.9005	0.0150	0.8970	0.0380
LL	4274.4950		4407.2970		5816.9220		5917.6170	
AIC	-8540.9905		-8806.5938		-11625.8442		-11827.2338	
BIC	-8519.3661		-8784.9694		-11604.2198		-11805.6094	
GARCH-t								
	BIST30		BIST100		USD		EURO	
Parameter	Value	Std	Value	Std	Value	Std	Value	Std
μ	0.0013	0.0000	0.0014	0.0000	-0.0001	0.0000	0.0000	0.0000
α_0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
α_1	0.1080	0.0340	0.1237	0.0420	0.1000	0.0170	0.1471	0.0390
β	0.8613	0.0500	0.8485	0.0510	0.8985	0.0170	0.8529	0.0400
d	8.0756	1.825	7.3133	1.7420	8.4006	1.2500	4.6384	0.2690
LL	4305.663		4446.708		5845.831		5949.975	
AIC	-8601.3251		-8883.4169		-11681.6623		-11889.9500	
BIC	-8574.2946		-8856.3854		-11654.6317		-11862.9194	

Table 4. Parameter estimates of GJR-n and GJR-t models and statistic tests

GJR-n								
	BIST30		BIST100		USD		EURO	
Parameter	Value	Std	Value	Std	Value	Std	Value	Std
μ	0.0010	0.0000	0.0014	0.0000	0.0000	4825134.081	-0.0001	0.001
α_0	0.0000	0.0000	0.0000	0.0000	0.0000	15973.427	0.0000	0.0000
α_1	0.0480	0.0015	0.0502	0.0140	0.2017	683963044.071	0.2889	0.0000
β	0.8501	0.0510	0.8487	0.0400	0.6650	675977556.066	0.5453	1.2460
γ	0.1255	0.0610	0.1490	0.0610	0.0000	5416656809.538	0.1305	1.9300
LL	4285.5600		4416.5190		5756.1670		5862.2690	
AIC	-8561.1200		-8823.0377		-11502.3339		-11714.5376	
BIC	-8534.0895		-8796.0071		-11475.3034		-11687.5071	
GJR-t								
	BIST30		BIST100		USD		EURO	
Parameter	Value	Std	Value	Std	Value	Std	Value	Std
μ	0.0010	0.0000	0.0011	0.0000	-0.0002	1406475.109	0.0000	28078131.946
α_0	0.0000	0.0000	0.0000	0.0000	0.0000	8879.443	0.0000	221202.851
α_1	0.0532	0.0170	0.0618	0.0180	0.3678	276992226.475	0.2539	29560250106.651
β	0.8371	0.0700	0.8198	0.0610	0.6226	426936136.846	0.7147	2212994007.407
γ	0.1294	0.0800	0.1291	0.0660	0.0000	2473886510.168	0.0003	47746730949.042
d	7.9437	1.5840	7.2247	1.2100	5.2103	5946045507.02	3.9473	13182520678.426
LL	4312.1270		4453.5100		5810.7600		5937.0560	
AIC	-8612.2540		-8895.0195		-11609.5192		-11862.1122	
BIC	-8579.8174		-8862.5829		-11577.0826		-11829.6756	

Table 5. Parameter estimates for static and dynamic copula families and model selection statistic

Static Copula	Parameter	GARCH-n	GARCH-t	GJR-n	GJR-t
Gaussian	ρ	0.0221	0.0297	0.0179	0.0360
	d	0.9731	0.9686	0.9664	0.9611
	LL	4351.2600	4329.1890	4055.8200	4347.8940
	AIC	-8704.5208	-8654.3780	-8107.6392	-8691.7872
	BIC	-8693.7086	-8643.5658	-8096.8270	-8680.9750
Student-t	d	6.5491	8.0109	23.9706	5.5313
	LL	4290.433	4243.688	3981.715	4330.352
	AIC	-8578.8664	-8485.9692	-7961.4291	-8658.7042
	BIC	-8573.4603	-8479.9692	-7956.0230	-8653.2981
Clayton	ω	0.85	0.85	0.85	0.85
	LL	3103.124	3103.277	2833.643	3101.575
	AIC	-6204.2483	-6204.5538	-5665.2859	-6201.1500
	BIC	-6198.8422	-6199.1477	-5659.8798	-6195.7439
SJC	λ	0.5353 0.5322	0.8109 0.8144	0.5348 0.5388	0.8299 0.8319
	LL	1413.471	2299.055	1443.225	2424.352
	AIC	-2822.9424	-4594.1105	-2882.4507	-4844.7030
	BIC	-2812.1302	-4583.2983	-2871.6385	-4833.8908

Dynamic Copula	Parameter	GARCH-n	GARCH-t	GJR-n	GJR-t
GDCC	α	0.0316	0.0297	0.0179	0.036
	β	0.9663	0.9686	0.9663	0.9611
	LL	4358.363	4329.189	4055.820	4347.894
	AIC	-8712.7252	-8654.3780	-8107.6391	-8691.7870
	BIC	-8701.9130	-8643.5658	-8096.8269	-8680.9748
tDCC	d	7.5376	9.5010	24.9955	5.9672
	α	0.028	0.0266	0.0194	0.0286
	β	0.9703	0.9718	0.9684	0.9688
	LL	4409.837	4365.207	4083.280	4418.826
	AIC	-8813.6744	-8724.4134	-8160.56	-8831.6529
tvC	BIC	-8797.4561	-8708.1951	-8144.3417	-8815.4346
	ω	12.8923	12.8913	12.8748	12.8905
	α	0.2197	0.2223	0.2343	0.2222
	β	12.7842	12.7839	12.704	12.7857
	LL	2430.023	2433.630	2403.856	2423.371
tvSJC	AIC	-4854.0459	-4861.2597	-4801.7120	-4840.7423
	BIC	-4837.8276	-4845.0414	-4785.4937	-4824.5240
	λ	9.1147 1.7757	-2.3995 1.5160	-0.4491 1.1956	1.4481 9.9382
	α	1.0295 8.4754	-8.3109 7.6800	-0.3626 1.2357	9.9486 9.6797
	β	3.228 3.9021	2.9388 5.0730	-0.4471 -0.3567	2.6305 3.0602
	LL	1966.988	1706.809	1431.022	1976.633
	AIC	-3921.9770	-3401.6179	-2850.0436	-3941.2664
	BIC	-3889.5404	-3369.1812	-2817.6070	-3908.8298

Table 6. The results of portfolio optimization based on CVaR

β	VaR	$CVaR$	x_1 -BIST30	x_2 -BIST100	x_3 -USD	x_4 -EURO
90%	0.0057	0.0102	0.3888	0.1552	0.2676	0.1884
95%	0.0084	0.0136	0.3760	0.1611	0.2715	0.1913
99%	0.0159	0.0233	0.3535	0.1486	0.2983	0.1996

5. CONCLUSION AND DISCUSSION

In this study, the dynamic Copula model was introduced and applied to four different sets of financial data (BIST30, BIST100, USD, EURO) in Turkey. In order to shape the model, first, the marginal distributions of the data series were determined as GJR-t, and by using identified marginal distributions, the copula model that established the dependency structure between the data series was selected as dynamic t (tDCC). By conducting the simulation, portfolio optimization was achieved based on CVaR risk measurement. Therefore, it can be inferred that investment in BIST30 is the best at each confidential level in accordance with the portfolio optimization results based on the minimization of CVaR. Other best investments are USD, EURO, and BIST100 respectively.

It is possible to add extreme value theory to the model due to the fat tail of financial data. Another possibility is to take into account the change points; we can offer different models for every period. Those will be topics of the future work.

REFERENCES

- [1] Markowitz, H. (1952) 'Portfolio Selection. Journal of Finance', vol. 25, pp. 77-91.
- [2] Rockafellar, R. T. and Ursayev, S. (2002) 'Optimization of conditional value at risk', Journal of Risk, vol. 2, pp. 21-40.
- [3] Sklar, A. (1959) 'Functions De Repartition An Dimensions At Leurs Marges', Publications De L'institut De Statistique De L'université De Paris, vol. 8, pp. 229-231.
- [4] He, H. and Li, P. (2011) 'Dynamic Asset Allocation Based on Copula and CVaR', IEEE.
- [5] Cherubini, U., Luciano, E. and Vecchiato, W. (2004) Copula Methods in Finance. John Wiley and Sons, New York.
- [6] Joe, H. and Xu, J.J. (1996) The estimation method of inference functions for margins for multivariate models. Technical Report 166. Department of Statistics, University of British Columbia.
- [7] Nelsen, R.B. (2006) An Introduction to Copulas, (second ed) Springer, New York.
- [8] Emrechts, P., McNeil, A. and Straumann D. (2002). 'Correlation and dependence in risk management: properties and pitfalls', In Risk Management Value at Risk and Beyond (Edited by M. Dempster), pp. 176-223, Cambridge University Press.
- [9] Rockinger, M. and Jondeau, E. (2006) 'The Copula-GARCH model of conditional dependencies: An international stock market application', Journal of International Money and Finance, vol. 25(3), pp. 827-853.
- [10] Wei, Y. and Zhang, S. (2007) 'Multivariate Copula-GARCH Model and Its Applications in Financial Risk Analysis [J].', Application of Statistics and Management, vol. 3, pp. 008.
- [11] Ozun, A., Cifter, A. (2007) 'Portfolio value-at-risk with time-varying copula: Evidence from the Americans', Marmara University. MPRA Paper No. 2711.
- [12] Huang, J. J., Lee, K. J., Liang, H. and Lin, W.F. (2009) 'Estimating value at risk of portfolio by conditional copula-GARCH method', Insurance: Mathematics and Economics, vol. 45, pp. 315- 324.
- [13] Wu, Z.X., Chen, M. and Ye, W.Y. (2006) 'Risk Analysis of Portfolio by Copula-GARCH', Journal of Systems Engineering Theory and Practice, vol. 2(8), pp. 45-52.
- [14] Wang, Z. R., Chen, X. H., Jin, Y. B., Zhou, Y.J. (2010) 'Estimating risk of foreign exchange portfolio: Using VaR and CVaR based on GARCH-EVT-Copula model', Physica A, vol. 389, pp. 4918-4928.

- [15] Patton, A.J. (2006) 'Modelling asymmetric exchange rate dependence', *International Economic Review*, vol. 47, pp. 527-556.
- [16] Bollerslev, T. (1986) 'Generalized autoregressive conditional heteroskedasticity', *Journal of econometrics*, vol. 31 (3), pp. 307-327.
- [17] Akaike, H. (1974) 'A new look at the statistical model identification', *Automatic Control, IEEE Transactions on*, vol. 19, pp. 716-723.
- [18] Schwarz, G. (1978) 'Estimating the dimension of a model', *Annals of Statistics*, vol. 6, pp. 461-464.
- [19] Cryer, C.D. and Chan, K.S. (2008). *Time Series Analysis with Applications in R*, (second ed.) Springer, New York.
- [20] <http://borsaistanbul.com/veriler/verileralt/hisse-senetleri-piyasasi-verileri/endeks-verileri> [02.08.2013].
- [21] <http://evds.tcmb.gov.tr/cbt.html> [02.08.2013].

Address: Ankara University, Faculty of Sciences, Department. of Statistics, Ankara, Kirikkale University, Faculty of Arts and Sciences, Department of Actuarial Science, Kirikkale, TURKEY
E-mail: acik@science.ankara.edu.tr, emel.kizilok@gmail.com

⁰Başlık:Bağımlı finansal varlıkların dinamik bağ ve CVaR'a dayalı portföy optimizasyonu ile modellenmesi

Anahtar Kelimeler:Dinamik Bağ, CVaR, portföy optimizasyonu