



Multiple Metastases of Observed Osteosarcoma Case in A Belgian Malinois Race Female Dog

Emin KARAKURT Hilmi NUHOĞLU Serpil DAĞ Enver BEYTUT

Kafkas University, Faculty of Veterinary Medicine, Department of Pathology, Kars, Turkey

Received: 09.04.2019

Accepted: 18.10.2019

ABSTRACT

In this study, we aimed to evaluate histopathologically the case of osteosarcoma detected in an Belgian Malinois race dog. The material of the study consisted of a 8.5-year-old Belgian Malinois race female dog weighing 30 kg. With the complaints of abdominal distention, difficulty in breathing, loss of appetite, fatigue, the dog brought to the Veterinary Faculty of Kafkas University Animal Hospital died on the way and was sent to the Department of Pathology for necropsy. In the follow-up, some of the tissue samples taken from the systemic necropsy animal were determined in the decalcified solution and another part in the buffered formaldehyde solution. Following routine procedures, 5 µm thick sections were taken from the prepared paraffin blocks for Hematoxylin Eosin staining. To determine the histopathological changes, the sections were examined by light microscope. In the light of the histopathological findings, osteosarcoma was diagnosed in the brain and lung metastatic foci.

Keywords: *Histopathology, Dog, Metastasis, Osteosarcoma*

öz

Belçika Malinois Irkı Dişi Bir Köpekte Gözlenen Osteosarkom Vakasına Ait Çoklu Metastazlar

Bu çalışmamızda, Belçika Malinois ırkı bir köpekte saptanan osteosarkom vakasının histopatolojik açıdan değerlendirilmesi amaçlanmıştır. Çalışmanın materyalini 8.5 yaşında 30 kg ağırlığında Belçika Malinois ırkı dişi bir köpek oluşturdu. Abdominal şişkinlik, solunum güçlüğü, iştahsızlık, halsizlik şikâyetleri ile Kafkas Üniversitesi Veteriner Fakültesi Hayvan hastanesine getirilen köpek yolda ölmüş ve nekropsisi yapılmak üzere Patoloji Anabilim Dalına gönderilmiştir. Takibinde sistemik nekropsisi gerçekleştirilen hayvandan alınan doku örneklerinin bir kısmı dekalsifiye solüsyonunda diğer bir kısmı ise tamponlu formaldehit solüsyonunda tespit edildi. Rutin işlemlerin ardından hazırlanan parafin bloklardan, Hematoksilen Eozin boyaması için 5 µm kalınlığında kesitler alındı. Histopatolojik değişikliklerin belirlenmesi amacıyla kesitler ışık mikroskopunda incelendi. Elde edilen histopatolojik bulgular ışığında beyin ve akciğerdeki metastatik odaklara osteosarkom tanısı konuldu.

Anahtar Kelimeler: *Histopatoloji, Köpek, Metastaz, Osteosarkom*

INTRODUCTION

Osteosarcoma is the most common primary bone tumor in dogs (Wehrle-Martinez et al. 2016). It constitutes 85% of skeletal malignancies and 4% of all malignancies in dogs (Wolfesberger et al. 2006; Milanta et al. 2012). This tumor with mesenchymal origin has very aggressive effects locally; It causes lysis or bone production; or causes both of these (McNeill et al. 2007; Petty et al. 2008). Metastasis is very common, continues usually subclinically and occurs early in tumor formation (Ohta et al. 2004; Fan et al. 2008; Shoeneman et al. 2012). Even osteosarcomas originating from long bones can metastasize to the lung even in the early period when the diagnosis of primary tumor cannot

be established (Sagartz et al. 1996). Lung metastasis occurs by hematogenous route and has been seen in 90% of all cases (Fieten et al. 2009; Piskun et al. 2011). Most of the dogs die due to lung metastasis (Gebhard et al. 2016). The prognosis of osteosarcoma is very poor, the recurrence rate is quite high; and less than 20% of the affected dogs do not live for more than 2 years (Wilson et al. 2008; Leonardo et al. 2018; Withers et al. 2019). Osteosarcoma is frequently observed in middle or old, male and large race dogs (Kirpensteijn et al. 2002; Hoenerhoff et al. 2004; Mullins et al. 2004; Selvarajah et al. 2012). Osteosarcomas are seen in high rate in races such as especially Doberman Pinscher, Golden Retriever, Great Dane, German Shepherd, St Bernard, Irish Setter, Boxer,

Rottweiler and Greyhound (McNeill et al. 2007; Cristo et al. 2017). 20-25% of skeletal osteosarcomas in dogs are caused by axial skeletal bones. Approximately 50% of these cases have axial skeletal osteosarcoma in head and in the other 50%; ribs, vertebrae and pelvis. Osteosarcoma observed in the head region is a rare tumor that is localized in the mandibula maxilla, skull and paranasal sinuses in dogs. The appendix skeleton is affected 3 to 4 times more than the axial skeleton (Nagamine et al. 2015; Murphy et al. 2017). Osteosarcoma is usually occurs in the metaphyseal region of the long bones (Sanches et al. 2009). The proximal humerus, distal radius, distal femur, proximal and distal tibia are the most common sites of appendicular osteosarcoma (Hoenerhoff et al. 2004). Amputation and chemotherapy are the best treatment options; untreated lesions; It has a tendency to progress rapidly and causes lameness and pathological fractures (Farese et al. 2004).

In this study, we aimed to evaluate histopathologically the case of osteosarcoma detected in a Belgian Malinois race dog.

CASE HISTORY

The material of the study consisted of an 8.5-year-old Belgian Malinois race female dog weighing 30 kg. With the complaints of abdominal distension, difficulty in breathing, loss of appetite and fatigue, the dog who was brought to the Veterinary Faculty Animal Hospital of Kafkas University died on the way and was sent to the Department of Pathology for necropsy.

In the follow-up, some of the tissue samples taken from the systemic necropsy animal were determined in decalcified solution and another part was detected in buffered formaldehyde solution. Following the routine procedures, 5 µm thick sections were taken from the paraffin blocks prepared for Hematoxylin&Eosin staining. To determine the histopathological changes, the sections were examined by light microscope. At the lower left part of the mandible, a mass of 4*4*3 cm fluctuated content was found. It was observed that the torsion around the long axis of the stomach was excessive and tympanic. Growing up in the spleen was seen. The examination of the intestine revealed that jejunum was hyperemic. The lungs were hyperemic and hard structures with a size of 1-3 mm were found in the parenchyma (Figure 1).




Figure 1. Lung tissue, hard structures with a size of 1-3 mm in the parenchyma (arrows)

The presence of white foci in the meninges was observed (Figure 2). The liver was mildly icteric. No other macroscopic findings were observed in the other organs and they showed normal anatomical structure. Although

the lung and brain metastasis sites were observed, primary tumor focus (anterior and posterior extremities, ribs and skull bones) could not be detected despite the detailed examinations.




Figure 2. Brain tissue, white foci in meninges (arrows)

Histopathological examination of tissue samples taken; degeneration of neurons, gliosis and osteosarcoma metastasis areas were observed in the brain. Dilatation of the sinusoids, dissociation in the remark cords, bilirubin pigment were common in hepatocytes. Hyperemia, atelectatic areas, spillage in bronchial and bronchiolar epithelium and anthracosis were observed in the lungs. Osteosarcoma metastasis sites were detected in the regions near the lung serosal surface (Figure 3).




Figure 3. Lung tissue, metastatic area (arrow), 500 µm , H&E

Purple-pink metastatic areas were found in the brain, similar to the foci in the lungs (Figure 4-5). Haemorrhagic areas of the spleen were found. Severe hyperemia in the jejunum, spillage in the mucosa epithelium and mononuclear cell infiltration in the lamina propria were determined.

Connective tissue increase in renal interstitium; hyaline cylinders in tubules; atrophy in glomeruli; edema fluid in the bowman capsule, andd chronic nonpurulent nephritis was detected. In the upper 1/3 of the lamina propria in the stomach lamina epithelials, necrosis was lost in severe hyperemia, mucosa and gland epithelium. It was determined that the mass taken under the mandible was cystic dilatation covered by a very folded epithelium. No histopathological findings were observed in the other organs and normal histological structure was observed.




Figure 4. Brain tissue (not decalcified), metastatic foci (arrows), 200 μ m, H&E

DISCUSSION and CONCLUSION

Osteosarcoma is often observed in males and large race dogs (Kirpensteijn et al. 2002; Hoenerhoff et al. 2004; Mullins et al. 2004; Selvarajah et al. 2012). The average age of the tumor is 7-8 years. Osteosarcomas in races such as Doberman, Pinscher, Golden Retriever, Great Dane, German Shepherd, St. Bernard, Irish Setter, Boxer, Rottweiler and Greyhounds are highly prevalent (McNeill et al. 2007; Cristo et al. 2017). There are only 5 studies in the Belgian Malinois in the diagnosis of osteosarcoma (Peterson et al. 2000; Selvarajah et al. 2009; Sivacolundhu et al. 2013, Moore et al. 2000, Pagano et al. 2016). In this study, we observed that a female dog of males having a males age of 8.5 years old with a weight of 30 kg was observed. The gender of the dog is different from the literature data indicating that this tumor is usually in male dogs. In this study, we observed that the osteosarcoma was detected in an 8.5-year-old-age Belgium Malinois race female dog weighing 30 kg; Although the age range and race data are consistent with the literature, being female of the dog is different from the literature data indicating that this tumor is usually in male dogs.

Osteosarcoma usually occurs in the metaphyseal region of the long bones (Sanches et al. 2009). The proximal humerus, distal radius, distal femur, proximal and distal tibia are the most common sites of appendicular osteosarcoma (Hoenerhoff et al. 2004). Cases originating from long bones in dogs usually cause lung, regional lymph nodes, liver and kidney organs (Kutsal et al. 2003; Murphy et al. 2017; Thompson and Dittmer 2017). It has rarely been reported to metastasize to the central nervous system, especially to the brain tissue (McNeill et al. 2007; Pazzi et al. 2013; Cristo et al. 2017). In our study, although there were metastases in the lung and brain similar to the literature data; In spite of the detailed examinations, primary tumor focus (anterior and posterior extremities, ribs and skull bones) could not be detected.

Based on the histopathological findings, osteosarcoma was diagnosed in the metastatic foci of the brain and lung tissue. In the literature search, it was observed that there were very few cases of osteosarcoma observed in the Belgian Malinois race. In addition, the material of the study material animal is female and the presence of metastatic foci in the brain, except for the lung tissue, makes the findings even more remarkable. Cerebral metastases of osteosarcoma cases observed in dogs are very rare and there are few literature studies on the subject. In Belgium




Figure 5. Brain tissue (decalcified), * means brain tissue, trabecular structures (arrows), 50 μ m, H&E

Malinois race dog, it is the first osteosarcoma study in which brain metastases have been shown. It is thought that this case report will make a significant contribution to the literature data.

REFERENCES

- Cristo TG, Vargas CB, Biezu G, Costa LS, Dal Pont TP, Kanamura CT, Salbego FZ, Traverso SD, Casagrande RA (2017). Metastatic osteosarcoma as a cause of hemorrhagic stroke in a dog. *Braz J Vet Pathol*, 10 (3), 105-110.
- Fan TM, Barger AM, Sprandel IT, Fredrickson RL (2008). Investigating TrkA expression in canine appendicular osteosarcoma. *J Vet Intern Med*, 22 (5), 1181-1188.
- Farese JP, Ashton J, Milner R, Ambrose LL, Van Gilder J (2004). The effect of the bisphosphonate alendronate on viability of canine osteosarcoma cells in vitro. *In Vitro Cell Dev Biol Anim*, 40 (3-4), 113-117.
- Fieten H, Spee B, Ijzer J, Kik MJ, Penning LC, Kirpensteijn J (2009). Expression of hepatocyte growth factor and the proto-oncogenic receptor c-Met in canine osteosarcoma. *Vet Pathol*, 46 (5), 869-877.
- Gebhard C, Fuchs-Baumgartinger A, Razzazi-Fazeli E, Miller I, Walter I (2016). Distribution and activity levels of matrix metalloproteinase 2 and 9 in canine and feline osteosarcoma. *Can J Vet Res*, 80 (1), 66-73.
- Hoenerhoff MJ, Kiupel M, Rosenstein D, Pool RR (2004). Multipotential Osteosarcoma with Various Mesenchymal Differentiations in a Young Dog. *Vet Pathol*, 41 (3), 264-268.
- Kirpensteijn J, Timmermans-Sprang EP, van Garderen E, Rutteman GR, Lantinga-van Leeuwen IS, Mol JA (2002). Growth hormone gene expression in canine normal growth plates and spontaneous osteosarcoma. *Mol Cell Endocrinol*, 197 (1-2), 179-185.
- Kutsal O, Kaya Ü, Vural S, Sağlam M (2003). Köpek ve Kedilerde 1986-2000 Yılları Arasında Ankara'da İncelenen Kemik Tümörleri. *Türk J Vet Anim Sci*, 27: 109-115.
- Leonardo L, Laura P, Serena BM (2018). miR-1 and miR-133b expression in canine osteosarcoma. *Res Vet Sci*, 117:133-137.
- McNeill CJ, Overley B, Shofer FS, Kent MS, Clifford CA, Samluk M, Haney S, Van Winkle TJ, Sorenmo KU (2007). Characterization of the biological behaviour of appendicular osteosarcoma in Rottweilers and a comparison with other breeds: a review of 258 dogs. *Vet Comp Oncol*, 5 (2), 90-98.
- Millanta F, Asproni P, Cancedda S, Vignoli M, Bacci B, Poli A (2012). Immunohistochemical expression of COX-2, mPGES and EP2 receptor in normal and reactive canine bone and in canine osteosarcoma. *J Comp Pathol*, 147 (2-3), 153-160.
- Moore GE, Mathey WS, Eggers JS, Estep JS (2000). Osteosarcoma in adjacent lumbar vertebrae in a dog. *J Am Vet Med Assoc*, 217 (7), 1038-1040.
- Mullins MN, Lana SE, Dernell WS, Ogilvie GK, Withrow SJ, Ehrhart EJ (2004). Cyclooxygenase-2 expression in canine appendicular osteosarcomas. *J Vet Intern Med*, 18 (6), 859-865.
- Murphy BG, Mok MY, York D, Rebhun R, Woolard KD, Hillman C, Dickinson P, Skorupski K (2017). Evaluation of P16 expression in canine appendicular osteosarcoma. *BMC Veterinary Research*, 13 (1), 189.

- Nagamine E, Hirayama K, Matsuda K, Okamoto M, Ohmachi T, Kadosawa T, Taniyama H (2015). Diversity of Histologic Patterns and Expression of Cytoskeletal Proteins in Canine Skeletal Osteosarcoma. *Vet Pathol*, 52 (5), 977-984.
- Ohta G, Sakai H, Kachi S, Hirata A, Yonemaru K, Kitajima A, Yanai T, Masegi T (2004). Assessment of proliferative potentials of canine osteosarcomas and chondrosarcomas by MIB-1 immunohistochemistry and bromodeoxyuridine incorporation. *J Comp Pathol*, 131 (1), 18-27.
- Pagano C, Boudreaux B, Shiomitsu K (2016). Safety And Toxicity Of An Accelerated Coarsely Fractionated Radiation Protocol For Treatment Of Appendicular Osteosarcoma In 14 Dogs: 10 Gy × 2 Fractions. *Vet Radiol Ultrasound*, 57 (5), 551-556.
- Pazzi P, Tompkins S, Kirberger RM (2013). Canine spirocercosis-associated extraskelatal osteosarcoma with central nervous system metastasis. *J S Afr Vet Assoc*, 84(1), E1-4.
- Peterson MR, Frommelt RA, Dunn DG (2000). A study of the lifetime occurrence of neoplasia and breed differences in a cohort of German Shepherd Dogs and Belgian Malinois military working dogs that died in 1992. *J Vet Intern Med*, 14 (2), 140-145.
- Petty JC, Lana SE, Thamm DH, Charles JB, Bachand AM, Bush JM, Ehrhart EJ (2008). *Vet Comp Oncol*, 6 (2), 133-140.
- Piskun CM, Muthuswamy A, Huelsmeyer MK, Thompson V, Stein TJ (2011). Wnt/ β -catenin expression does not correlate with serum alkaline phosphatase concentration in canine osteosarcoma patients. *PLoS One*, 6 (10), e26106.
- Sagartz JE, Bodley WL, Gamblin RM, Couto CG, Tierney LA, Capen CC (1996). p53 tumor suppressor protein overexpression in osteogenic tumors of dogs. *Vet Pathol*, 33 (2), 213-221.
- Sanches DS, Pires CG, Fukumasu H, Cogliati B, Matsuzaki P, Chaible LM, Torres LN, Ferrigno CR, Dagli ML (2009). Expression of connexins in normal and neoplastic canine bone tissue. *Vet Pathol*, 46 (5), 846-859.
- Selvarajah GT, Verheije MH, Kik M, Slob A, Rottier PJ, Mol JA, Kirpensteijn J (2012). Expression of epidermal growth factor receptor in canine osteosarcoma: association with clinicopathological parameters and prognosis. *Vet J*, 193 (2), 412-419.
- Shoeneman JK, Ehrhart EJ, Eickhoff JC, Charles JB, Powers BE, Thamm DH (2012). Expression and function of survivin in canine osteosarcoma. *Cancer Res*, 72 (1), 249-259.
- Sivacolundhu RK, Runge JJ, Donovan TA, Barber LG, Saba CF, Clifford CA, de Lorimier LP, Atwater SW, DiBernardi L, Freeman KP, Bergman PJ (2013). Ulnar osteosarcoma in dogs: 30 cases (1992-2008). *J Am Vet Med Assoc*, 243 (1), 96-101.
- Thompson KG, Dittmer KE (2017). Tumors of Bone. In: *Tumors of Domestic Animals*, Moulton JE (Ed), 356-424, Wiley Blackwell, Iowa.
- Wehrle-Martinez AS, Dittmer KE, Aberdein D, Thompson KG (2016). Osteocalcin and Osteonectin Expression in Canine Osteosarcoma. *Vet Pathol*, 53 (4), 781-787.
- Wilson H, Huelsmeyer M, Chun R, Young KM, Friedrichs K, Argyle DJ (2008). Isolation and characterisation of cancer stem cells from canine osteosarcoma. *Vet J*, 175 (1), 69-75.
- Withers SS, Skorupski KA, York D, Choi JW, Woolard KD, Laufer-Amorim R, Sparger EE, Rodriguez CO, McSorley SJ, Monjazebe AM, Murphy WJ, Canter RJ, Rebhun RB (2019). Association of macrophage and lymphocyte infiltration with outcome in canine osteosarcoma. *Vet Comp Oncol*, 17 (1), 49-60.
- Wolfesberger B, Walter I, Hoelzl C, Thalhammer JG, Egerbacher M (2006). Antineoplastic effect of the cyclooxygenase inhibitor meloxicam on canine osteosarcoma cells. *Res Vet Sci*, 80 (3), 308-316.