

# Original Article

Ulutas Med J 2015;1(4):100-103  
DOI: 10.5455/umj.20151225044952  
ISSN:2149-0430 eISSN: 2149-388X



## Right Ventricle Diastolic Function in High-Normal and Mildly Elevated Arterial Blood Pressure

Noha Hassanin Hanboly

Faculty of Medicine, Cardiovascular Department, Cairo University, Egypt

**Background:** Many studies evaluated left ventricle (LV) function in mildly increased arterial blood pressure, however right ventricle (RV) had received scarce attention due to the complex shape and orientation. The aim of this study was to evaluate RV diastolic dysfunction using pulsed and tissue Doppler echocardiography in never-treated patients with blood pressure (BP) values varying from the optimal to the mild hypertensive range.

**Method:** The study group consisted of 57 patients with high-normal and mildly increased arterial BP. Twenty healthy age and sex matched subjects were recruited as controls. Conventional echocardiography was done to study LV dimensions and systolic function. Pulsed Doppler echocardiography was used to measure peak early and late tricuspid diastolic filling velocities. Tissue Doppler imaging was done to determine RV annular velocities.

**Results:** Right ventricle hypertrophy was found in 28% of patients with newly diagnosed systemic hypertension (n:16). Tricuspid E/A was 1.3 and 0.9 in control and hypertensive groups respectively. Tissue Doppler tricuspid E'/A' was significantly decreased in hypertensive group ( $0.8 \pm 0.2$  versus  $1.2 \pm 0.1$  in control,  $p < 0.001$ ).

**Conclusion:** right ventricle diastolic function deteriorates in response to slightly increased systemic blood pressure. Right ventricular diastolic dysfunction may be an early clue to hypertensive heart disease.

**Keywords:** Diastolic dysfunction, Right ventricle, Tissue doppler echocardiography.

### Introduction

Arterial systemic hypertension may determine impairment of both left ventricle (LV) diastolic and systolic functions, due to increased after load (1). Also the right ventricle (RV) might be involved in this process by structural and functional abnormalities (2).

Left ventricular hypertrophy is often associated with hypertension and is an adaptive mechanism to maintain or normalize wall stress, sometimes at the expense of diastolic and long-axis systolic function (3). This adaptive mechanism is associated with changes in left ventricular parameters and in chamber dimensions, geometry and function. These changes are progressive and can ultimately lead to heart failure with systolic

and/or diastolic dysfunction (4). Human heart functions as a unit that includes the right ventricle. The spectrum of changes in structure, function and shape of the left ventricle ultimately has an effect on the structure and function of the RV (5).

Only a few publications concern echocardiographic assessment of the RV diastolic function in systemic hypertension, which is all more surprising as physical examination often demonstrates the features of RV pathology and dysfunction in hypertensive patients (6). Purpose of the study was assessment of RV diastolic function in patients with untreated systemic hypertension using conventional and doppler echocardiography.

**Corresponding Author:** Noha Hassanin Hanboly. Faculty of Medicine, Cardiovascular Department, Cairo University, Egypt.

**E-mail:** Noha.Ali@kasralainy.edu.eg

**Received:** Nov 30, 2015 **Accepted:** Dec 25, 2015

**Published:** December 28, 2015

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

The Ulutas Medical Journal © 2015  
Available at <http://ulutasmedicaljournal.com>

## Study Design

The study was carried out at the echocardiography laboratory of cardiovascular department at Cairo University hospitals. The Research Ethics Committee of the hospital reviewed and approved the study protocol. Fifty seven adult participants aged  $54.8 \pm 14$  years old with high-normal and mildly increased arterial blood pressure never treated with antihypertensive medications were compared with 20 age and gender matched control subjects. Patient written consent was given by all the participants. Hypertension was defined as persistent elevation of blood pressure above 140/90 mmHg on two or more occasions with the patient in a sitting position for at least five minutes (7). Patients were subjected to history taking, physical examination and simple clinical investigations. Demographic parameters, including age, gender and associated symptoms were obtained. Subjects with ischemic heart disease, valvular heart disease and cor pulmonale were all excluded.

## Echocardiography

All subjects had echocardiography performed with the use of a Philips IE 33 (Andover, MA, USA) Ultrasound according to recommendations of the American Society of Echocardiography (8). Left ventricle structure, dimensions and functions were studied. Right ventricle wall thickness was measured in diastole from the sub costal views wall and wall thickness  $\geq 5$  mm indicates RV hypertrophy (9). Right ventricle diastolic function was assessed by recording tricuspid flow with standard pulsed wave Doppler and by measurements of early diastolic peak flow velocity (E), late diastolic peak flow velocity (A) and the ratio of early-to-late flow velocity peaks (E/A ratio).

## Tissue Doppler imaging (TDI)

In the apical four-chamber view, the sample volume was placed at the lateral tricuspid annulus to study early diastolic (E') and late diastolic (A') velocities. The use of E':A' ratio is one of the recommended indices approved by the American Society of Echocardiography for assessing right ventricle diastolic dysfunction (8, 10).

## Statistical analysis

All statistical calculations were done using computer program SPSS (Statistical Package for the Social Science, SPSS Inc., Chicago, IL, USA) release 15 for Microsoft Windows (2006). All Data were statistically described in terms of mean  $\pm$  standard deviation ( $\pm$  SD) Comparison of numerical variables between the study groups was done using Student t test for independent samples and one way analysis of variance (ANOVA) test when comparing

3 group. P-value less than 0.05 was considered statistically significant.

## Results

Fifty seven participants aged  $54.8 \pm 14$  years old (40 females and 17 males) with high-normal and mildly increased arterial blood pressure never treated with antihypertensive medications were compared with 20 age and gender matched control subjects free of any cardiac diseases. Systolic (SBP) and diastolic blood pressures (DBP) were significantly higher in hypertensive subjects compared to the controls ( $143.7 \pm 2.7$  and  $93.2 \pm 5.1$  mmHg vs  $119.4 \pm 8.1$  and  $72.3 \pm 6.9$  mm hg, p-value  $< 0.001$  for both)(Table-1).

**Table-1:** Demographic-clinical baseline data of study population

	Hypertensive group	Control group	P
Age, years	$54.8 \pm 14$	$54.2 \pm 17.2$	0.879
Body mass index, kg/m <sup>2</sup>	$33.3 \pm 7.4$	$25.9 \pm 7.2$	0.001*
Waist circumf. (WC), cm	$115 \pm 14.1$	$90.6 \pm 11.8$	0.001*
Systolic blood pressure, mmHg	$143.7 \pm 2.7$	$119.4 \pm 8.1$	$< 0.001^*$
Diastolic blood pressure, mmHg	$93.2 \pm 5.1$	$72.3 \pm 6.9$	$< 0.001^*$


(\* ) denotes statistically significant

## Conventional Echocardiography

Interventricular septum thickness and right ventricle free wall thickness were significantly increased in patients with mild untreated systemic hypertension ( $1.2 \pm 0.2$  and  $0.5 \pm 0.1$  vs.  $0.8 \pm 0.1$  cm and  $0.28 \pm 0.1$  cm among control group,  $P < 0.001$  for both).

## Pulsed and tissue Doppler imaging

Mitral E/A ratio  $1.2 \pm 0.1$  and  $0.7 \pm 0.1$  in the control and hypertensive group respectively (P value  $< 0.001$ ). While, tricuspid E/A was 1.3 and 0.9 in control and hypertensive groups respectively. Tissue Doppler tricuspid E' and E'/A' values were significantly decreased in hypertensive group ( $12 \pm 3.1$  cm/s and  $0.8 \pm 0.2$  vs  $15.3 \pm 1.3$  cm/s and  $1.2 \pm 0.1$  in control group, p  $< 0.001$  for both) (Figure-1). Positive significant correlation was found between mitral and tricuspid E/A ratio (p  $< 0.001$ , r: 0.842). Univariate regression analysis revealed that body mass index was found to be significant predictor of RV diastolic dysfunction (p:0.023).



**Figure-1:** Tissue Doppler imaging of right ventricle lateral tricuspid annulus in a normal subject E'/A' ratio is >1(a) and in a hypertensive patient E'/A' ratio <1(b).

## Discussion

It was found that in middle-aged men central distribution of body fat is associated with increased BP, independently of body mass index and insulin resistance, thus suggesting a key role of central adiposity in the full expression of the “metabolic syndrome (11).

The current study showed that body mass indices and waist circumferences were increased in the hypertensive group. Moreover, body mass index was found to be significant predictor of RV diastolic dysfunction (RVDD). This is concordant with other studies that found that mitral annular E':A' ratio and body mass index were the only predictors of RVDD (10). Concordant with other studies (12), the present study used the ratio E':A' ratio alone to assess RVDD. However, this index was approved by the American Society of Echocardiography to assess the global RV diastolic function. Positive significant correlation was found between mitral and tricuspid E/A ratio ( $p < 0.001$ ,  $r = 0.842$ ) this is in accordance with other study that assessed RV DD in 100 adult Nigerian hypertensive subjects (13).

Concordant with some studies that found that the prevalence of RV hypertrophy consistently varied among studies (17.0-80.0%) with an average of 28.6% in the pooled population (14), the current study found that RV hypertrophy was prevalent in 28% of the high-normal and mildly increased blood pressure group. It was found that RV diastolic and systolic function deteriorates in response to slightly increased systemic BP (15). It was found that increased pulmonary vascular resistance in systemic hypertension results from the activation of vasoconstriction pathways in both systemic and

pulmonary vascular beds which may explain right ventricular hypertrophy from mild to moderate, uncomplicated hypertensive patients (16).

Moreover, combination of increased thickness of the posterior wall, inter ventricular septum dimensions and right ventricular wall thickness will ultimately lead to at least progressive reduction in the right ventricular end-diastolic dimensions before progressive dilatation may occur in the right heart (17).

A magnetic resonance imaging study to assess RV remodeling in systemic hypertension revealed that that systemic hypertension was associated with concentric RV remodeling and impaired diastolic function, confirming that the unstressed ventricle is not immune to the effects of systemic hypertension (18).

## Conclusion

In arterial hypertension, increased systemic after load is transferred to RV, a structure not directly exposed to systemic BP but the inter ventricular septum is likely the candidate for that role. Right ventricle diastolic function could provide novel insights in RV function impairment in high-normal and mildly increased arterial blood pressure. Moreover, RV diastolic dysfunction may be an early clue to hypertensive heart disease.

## Conflict of Interest & Acknowledgement

All authors declared no conflict of interest. The assistance of staff members at echocardiography laboratory cardiovascular department Cairo University is greatly appreciated. No external source of funding was used for the current study.

## Reference

- Galderisi M, Petrocelli A, Alfieri A, Garofalo M, de Divitiis O. Impact of ambulatory blood pressure on left ventricular diastolic function. *Am J Cardiology* 1996, 77: 597-601.
- Myslinski W, Mosiewicz J, Ryczak E, Barud W, Bilan A, Palusinski R and Hanzlik J. Right ventricular function in systemic hypertension. *J Hum Hypertension* 1998, 12: 149-155.
- Verdecchia P, Schillaci G, Borgioni C, et al. Prognostic value of left ventricular mass and geometry in systemic hypertension with left ventricular hypertrophy. *Am J Cardiology* 1996, 78: 197-202.
- Janicki J. Influence of the pericardium and ventricular interdependence on left ventricular diastolic and systolic function in patients with heart failure. *Circulation* 1990, 81(suppl III): III-15-III-20.
- Clyne C, Alpert J, Benotti J. Interdependence of the left and right ventricles in health and disease. *Am Heart J* 1989, 117: 1366-1373.
- Cittadini A, Fazio S, Stromer H et al. Optimal determination of right ventricular filling dynamics in systemic hypertension. *Am Heart J* 1995, 130:1074-1082.

7. Paul A, Suzanne O, Barry L, William C, Cheryl D, Joel H, Daniel T, et al. Report From the Panel Members Appointed to the Eighth Joint National Committee (JNC). JAMA 2014, 311:507-520.
8. Lawrence G, Wyman W, Jonathan A, Lanqi H, Mark D, Krishnaswamy C, Scott D et al. Guidelines for the cardiography Assessment of the right heart in adults. J Am Soc cardiography 2010, 23:685-713.
9. Roberto M, Luigi P, Victor M, Jonathan A, Anderson A, Laura E, Frank A et al. Recommendations for Cardiac Chamber Quantification by Echocardiography in Adults: An Update from the American Society of Echocardiography and the European Association of Cardiovascular imaging. J Am Soc Echocardiogr 2015; 28:1-39.
10. Cicala S, Galderisi M, Caso P, Petrocelli A, D' Errico A, de Divitiis O et al. Right ventricular diastolic dysfunction in arterial systemic hypertension: analysis by pulsed tissue Doppler. Eur J Echocardiography 2002; 3:135-142.
11. Alfonso S, Francesco P, Gianvincenzo B. The Relationship of waist circumference to blood pressure. AJH 2002,15:780-786.
12. Kamilu M, Hadiza S, Mohammed N. Right ventricular dysfunction in a hypertensive population stratified by patterns of left ventricular geometry. Cardiovascular J Africa 2012; 23: 478-482.
13. Akintunde A, Akinwusi P, Familoni O and Familoni O. Effect of systemic hypertension on right ventricular morphology and function: an echocardiography study. Cardiovasc J Afr. 2010 Oct; 21(5): 252-256.
14. Cuspidi C, Sala C, Muiesan M, et al. Right ventricular hypertrophy in systemic hypertension: an updated review of clinical studies. J Hypertension 2013, 31:858-65.
15. Pedrinelli R, Canale M, Giannini C, Talini E, Penno G, Dell'Omo G, Di Bello V. Right ventricular dysfunction in early systemic hypertension: a tissue Doppler imaging study in patients with high-normal and mildly increased arterial blood pressure. J Hypertension 2010, 28:615-21.
16. Vriz O, Argiento P, D'Alto M, Ferrara F, Vanderpool R, Naeije R, Bossone E. Increased pulmonary vascular resistance in early stage systemic hypertension: a resting and exercise stress echocardiography study. Can J Cardiol 2015; 4:537-43.
17. Nunez B, Amodeo C, Garavaglia G, Schmiedt R, Frohlich E. Biventricular cardiac hypertrophy in essential hypertension. Am Heart J 1987;114:813-818.
18. Todiere G, et al. Right ventricular remodeling in systemic hypertension:a cardiac MRI study. Heart 2011, 97:1257-61.

**DOI:** [dx.doi.org/10.5455/umj.20151225044952](https://doi.org/10.5455/umj.20151225044952)

**Cite article as:** Noha Hassanin Hanboly. Right ventricle diastolic function in high-normal and mildly elevated arterial blood pressure. Ulutas Med J. 2015;1(4):100-103.

**Submit your next manuscript to The Ulutas Medical Journal and take full advantage of;**

- Convenient online submission
- Thorough peer review, **Fast Response**
- No space constraints or color figure charges
- **Immediate publication** on acceptance
- Inclusion in **Scopemed** and **Google Scholar**
- Research which is freely available for redistribution

**Submit your manuscript at:**

<http://ulutasmedicaljournal.com>



Ulutas Med J \* 2015;1(4):100-103