

# Case Report

Ulutas Med J 2015;1(4):119-121  
DOI: 10.5455/umj.20151110010021  
ISSN:2149-0430 eISSN: 2149-388X



## Suicidal Death from Pseudoephedrine Sulfate Overdose: A Case Report

Semih Petekkaya<sup>1</sup> Nusret Ayaz<sup>2</sup> Mustafa Doğan<sup>2</sup> Mucahit Oruc<sup>2</sup> Bedirhan Sezer Oner<sup>1</sup>  
Cihan Gokturk<sup>1</sup> Ahmet Celebi<sup>2</sup> Adalet Eda Budak<sup>1</sup> Ozcan Soylu<sup>1</sup> Osman Celbis<sup>2</sup>

<sup>1</sup> Council of Forensic Mortuary Department, Malatya, Turkey <sup>2</sup> Department of Forensic Medicine, Inonu University, Malatya, Turkey

**Background:** Upper respiratory tract infections are common in the community and pseudoephedrine preparations are widely used in the treatment as a decongestant drug. However, pseudoephedrine has serious side effects on the central nervous system and the cardiovascular system. Although cases of infant death due to overdose have been reported in literature, pseudoephedrine toxicity has not been observed in adults as a cause of suicide in literature.

**Case presentation:** The case is an 18-year-female committed suicide with pseudoephedrine sulfate preparation.

**Conclusion:** Care and attention must be paid as pseudoephedrine preparations are easily accessible, widely used and an overdose can cause death in adults.

**Keywords:** Pseudoephedrine sulfate, suicide, intoxication, death

### Introduction

Pseudoephedrine sulfate is a sympathomimetic agent stimulating Alpha and Beta-2 adrenergic receptors, which is widely used in upper respiratory tract infections. With the alpha adrenergic effect, there is a decrease in the findings of oedema and inflammation in the mucosa caused by vasoconstriction in the smooth muscles in the veins in the respiratory tract mucosa. The bronchodilator effect is shown by stimulation of the beta 2 adrenergic receptors in the bronchial smooth muscles (1). Daily oral doses are used as 60mg x 4/day for adults and a total of 4mg/kg/day as 4 equal doses for children (2, 3, 4). After oral intake, it is quickly and almost all absorbed from the digestive canal. After 3-6 hours, a 120mg dose reaches peak concentration of 265-314ng/ml. Less than 1% of the absorbed amount is transformed to an inactive metabolite with N-demethylation in the liver and 55-96% of the administered dose is eliminated from the body without

any change together with the urine. The elimination half-life has been reported as 4.3 - 8 hours (5).

In this paper, the clinical, autopsy, histopathology and toxicology findings are presented of an adult female who committed suicide through an overdose of pseudoephedrine. To the best of our knowledge, there is no similar case in literature.

### Case Report

An 18-year old female was brought to the emergency department approximately 90 mins after attempting suicide by taking 12-13 tablets each containing 120 mg pseudoephedrine sulfate and 5 mg loratadine agent. Stomach lavage was applied and the patient was admitted to the intensive care unit for monitoring and treatment. The vital signs of patient were stable up to 5 hours, then agitation developed, aggressive behaviour, tachycardia

**Corresponding Author:** Nusret Ayaz; Department of Forensic Medicine, Inonu University, Malatya, Turkey

**E-mail:** nusretayaz@gmail.com

**Received:** Oct 26, 2015 **Accepted:** Nov 10, 2015

**Published:** December 28, 2015

*This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.*

**The Ulutas Medical Journal** © 2015  
Available at <http://ulutasmedicaljournal.com>

(178 bpm), hypotension (90/50 mmHg), fever (39°C) and clouded consciousness, so she was intubated. In blood gas sample which was taken, pH was determined as 7.01, HCO as 12, and PCO<sub>2</sub> as 64. The necessary treatments were organised, transfer for hemodia filtration was planned, but 4 hours after the onset of the symptoms, the patient suffered cardiac arrest and died.



**Fig-1.** 150 ml serous fluid (Fig 1) in the pericardium



**Fig-2.** Widespread subpericardial and parenchymal bleeding



**Fig-3.** Beige-coloured solid particles in the stomach contents

In the autopsy, external examination revealed swelling on the left side of the lower lip and a 0.2 cm area of ecchymosis abrasion on the lower part, and needle marks on the back of the hand, the wrist, inside the elbow and in the left groin area. When the head cavity was opened, areas of petechiae under the skin with hair and a

hyperemic appearance of the brain, brain stem and cerebellum were determined. In chest cavity examination there was 150 ml serous fluid (Fig-1) in the pericardium and areas of widespread subpericardial and parenchymal bleeding (Fig-2) were seen. On the surface of both lungs, areas of petechial bleeding were seen. In the examination of the abdominal cavity, 50 ml dark-coloured fluid in the stomach contents was seen to contain beige-coloured solid particles (Fig-3). Petechial bleeding was seen in the superior anterior section of the liver.

For histopathological evaluation, samples were taken from the heart, brain, kidneys, liver, lungs, pancreas, oesophagus and stomach and placed in 10% formaldehyde and sent to the pathology laboratory. Following routine tissue testing, the samples taken from the tissues were stained with hematoxylin-eosin (H-E) and evaluated with a light microscope. In the microscopic examination, focal points of bleeding were determined in the brain stem and pons parenchyma, widespread fresh bleeding in the parenchyma of the heart, no pathology of note was observed in the coronary arteries and autolysis was determined in the stomach mucosa.

Samples of the liver, kidney, stomach contents, fat tissue, bile fluid and blood were taken and sent for systematic toxicology laboratory evaluation. In the examination of the samples with a Gas Chromatography /Mass Spectrometer (GC/MS) device, there was seen to be pseudoephedrine agent substance in the blood, organ parts and stomach contents. In the examination of blood made with Liquid Chromatography /Mass Spectrometer (LC/MS) device, the pseudoephedrine value was determined as 13638 ng/ml. The cause of death was reported as respiratory and cardiovascular effects due to high dosage pseudoephedrine toxicity.

## Discussion

Pseudoephedrine demonstrates an effect on both alpha and beta adrenergic receptors. In cases of overdose, by affecting these receptors, effects are seen which may range from depression of the central nervous system (*sedation, apnea, loss of attention, cyanosis, coma, cardiovascular collapse*) to stimulation (*insomnia, agitation, hallucinations, tremor or convulsions*) from cardiovascular effects (*tachycardia, hypo-hypertension, cardiac arrhythmia, myocardial infarctus*), hyperthermia and death (6, 7, 9, 10).

Pseudoephedrine has been reported to have effects on cardiovascular system of arrhythmia, palpitation, increased systolic blood pressure, tachycardia, cardiovascular collapse, paroxymal supra ventricular arrhythmia and

to cause myocardial infarctus without any pathological change in the coronary artery structure (9, 10). In the case presented here, tachycardia and hypotension were reported during the clinical observation period. In the macroscopic findings of the autopsy, there were areas of widespread subepicardial and parenchymal bleeding in the heart and in the microscopic examination, areas of widespread fresh bleeding were seen in the parenchyma of the heart without any remarkable pathology in the coronary arteries. These findings were thought to have occurred as a result of the catecholaminergic effect associated with the sympathetic system stimulation of pseudoephedrine.

By affecting the adrenergic receptors in the central nervous system, pseudoephedrine enables the expression of neuradrenalin from the neurons. A stimulant effect is shown in the central nervous system and there is an increase in sympathetic tonus. A treatment dosage may lead to hallucinations, behavioural changes and seizures while a high dosage causes agitation, psychosis, paranoia and intercranial haemorrhage (4, 11). In the case, agitation and aggressive behaviour were reported in the clinical observation period. In the macroscopic findings of the autopsy there was a hyperemic appearance of the brain, cerebellum and brain stem and in the microscope examination, focal points of bleeding were determined in the brain stem and pons parenchyma were seen.

Toxic dose effects of pseudoephedrine have been reported following the intake of 4-5 times more than the therapeutic dose (180-360mg)(7). In blood examination in the current case with GC/MS and LC/MS devices, the pseudoephedrine value was determined as 13638 ng/ml, which was the level of a toxic dose. Clinical, microscopic and macroscopic findings were seen to be consistent with the side-effects of pseudoephedrine and effects associated with a high dose intake as described in literature (3, 4, 8, 9, 10, 11).

## Conclusion

Cases of pseudo ephedrine-related deaths have generally been reported as occurring in infants when the treatment dose has been exceeded. Although there have been several reported cases of side-effects seen in adults taking pseudoephedrine, to the best of our knowledge there have been no reported cases of the death of an adult as a result of a high dose of pseudoephedrine taken with the aim of suicide. Therefore, this can be considered as of importance as the first reported case of pseudoephedrine-related suicide.

## Conflict of Interest & Acknowledgement

The authors declare that no conflict of interest exists in publishing this article. This case has been presented as poster at the 1st International Congress and Workshop of Forensic Toxicology, at the date of 29<sup>th</sup> and 30<sup>th</sup> November 2014 in Ankara, Turkey.

## Reference

1. Kayaalp O, Akılcı Tedavi Yönünden Tıbbi Farmakoloji 2012; 2 (13): 1022-1023. Ankara. Pelikan Kitabevi.
2. Gülhan B, Bayrakçı B, Babaoglu M Ö, Bal B, Beken S. Biphasic reatin kinase elevation in pseudoephedrine overdose. *Br J Clin Pharmacol* 2008; 67:139-140.
3. Soyer T, Göl IH, Eroğlu F, Cetin A. Acute urinary retention due to pseudoephedrine hydrochloride in a 3-year-old child. *Turk J Pediatr.* 2008; 50:98-100.
4. Roberge RJ, Hirani KH, Rowland PL 3rd, Berkeley R, Krenzelok EP. Dextromethorphan- And Pseudoephedrine-Induced Agitated Psychosis And Ataxia. *J Emerg Med.* 1999 Mar-Apr;17:285-8.
5. Brunton L, Chabner B, Knollman B, eds. Goodman& Gilman's The Pharmacological Basis Of Therapeutics. 12th edn. China: The McGraw Hill Company; 2011.
6. Marinetti L, Lehman L, Casto B, Harshbarger K, Kubiczek P, Davis J, Over-the-Counter Cold Medications – Postmortem Findings in Infants and the Relationship to Cause of Death. *Journal of Analytical Toxicology.* 2005; 29:738-743.
7. Olson K. R, Lange Zehirlenmeler & ilaç Aşırı Dozu. Istanbul: Nobel Tıp Kitabevi. 2012.
8. Kutluhan S. Özen A. Gökalp O. Gültekin F. Psödoefedrin ile Tetiklenen Epileptik Nöbet. *Türk Nöroloji Dergisi.* 2008;14:357-359.
9. Aypak C, Türedi Ö, Yüce A, Görpelioğlu S. A Serious Adverse Effect of Pseudoephedrine Used For Common Cold Treatment : Ventricular Arrhythmia. *Cukurova Med J* 2013; 38: 506-510.
10. Tahmaz M, Kumbasar B, Ozturk S, Kazancıoğlu R. Pseudoephedrine-induced Ventricular Tachycardia. *Eur J Gen Med* 2013; 10:77-80.
11. Loizou LA, Hamilton JG, Tsementzis SA. Intracranial hemorrhage in association with pseudoephedrine overdose (letter). *J Neurol, Neurosurg Psychiatr* 1982;45:471-2.

DOI: [dx.doi.org/10.5455/umj.20151110010021](https://doi.org/10.5455/umj.20151110010021)

**Cite article as:** Petekkaya S, Ayaz N, Dogan M, Oruc M, Oner BS, Gokturk C, Celebi A, Budak AE, Soyulu O, Celbis O. Suicidal Death from Pseudoephedrine Sulfate Overdose: A Case Report. *Ulutas Med J.* 2015;1(4):119-121