

ISSN: 2149-6307 Lycia Press LONDON UK





International Journal of Natural Science and Discovery **Open Access Scientific Journal** 2016, Vol.2, No.4

www.natscidiscovery.com

Natural Science and Discovery (www.natscidiscovery.com) is an international open access, peer-reviewed scientific research journal that provides rapid publication of articles in all disciplines of Natural science such as Physics, Mathematics, Chemistry, Biology, Engineering, Nanotechnology

The policy of top priority of NSD is to put forward and highlight medical innovations and inspiring patents

NSD offers an exceptionally fast publication schedule including prompt peer-review by the experts in the field and immediate publication upon acceptance.

The editorial board aims at reviewing the submitted articles as fast as possible and promptly including them in the forthcoming issues.

This journal is published under ethical publishing policy of international scientific ethical and publication rules. NSD supports the Open Access Initiative. Abstracts and full texts (HTML and PDF format) of all articles published by NSD are freely accessible to Researchers, Scientist, the Public and other users to view, download and share, immediately upon publication

Honorary Editors

Prof. Dr. Gustavo Barbosa-CÁNOVAS, Washington State University, USA Prof. Dr. Peyami BATTAL, Yuzuncu Yil University, TR

Natural Science and Discovery is an international open access, peer-reviewed scientific research journal. All concession and copyrights belonging to Lycia Press as Founder

ISSN: 2148-6832 (Print) E-ISSN: 2149-6307 (Online) Category: Multi Disciplinary Life Science Journal Abbreviated key title: Nat. Sci. Discov. Frequency: Sessional (March, June, September, December) Review System: Blind Peer Review Circulation: Globally, Online, Printed Article Processing Charge (APC): US\$ 100 Licensing: CC-BY-NC 4.0 International License Environmental Editor-in-Chief: Christopher E. SCHMİTT (Ph.D.) Cardiovascular Research Institute | San Francisco, CA, USA Deputy Editor: Halit Demir (Prof.) YYU, Faculty of Science Dept. of Biochemistry, Manisa, TR Deputy Editor: Zafer Akan (Assoc. Prof.) 3rd Floor 86 - 90 Paul Street, EC2A 4NE, London, UK Established: 30.04.2015 Web address: www.medscidiscovery.com; http://dergipark.ulakbim.gov.tr/msd E-mail : editor [at] medscidiscovery.com

Design and preparation of PDFs, Language editing, Web site design, Graphical design Services of international Journal of Medical Science and Discovery has been contracted with Lycia Press LONDON, UK (as Publisher), by the MSD Board of Directors

Publisher: Lycia Press Inc. Address: 3rd Floor 86 - 90 Paul Street, EC2A 4NE, London, UK Web address: www.lycians.com Phone : +44 0 203 322 3097 E-mail : office [at] lycians.com E-mail : info [at] lycians.com

Editorial Board

| Ph.D. | Christopher E. Schmitt | UCSF Cardiovascular Research Inst. San Francisco, CA, USA (Editor in Chief, |
|-------------------|------------------------|---|
| Prof. Dr. | Arash KHAKI | Islamic Azad University, Iran |
| Ph.D. | Muhammet Burak BATIR | Celal Bayar University,TR |
| Asist. Prof. Dr. | Ozdemirhan SERCIN | University Libre de Bruxelles, Belgium |
| Prof. Dr. | Nevin ÇANKAYA | USAK University TR |
| Asist. Prof. Dr. | Serife Kaymaksiz | Helmholtz-Institute Muenster, Germany |
| Ph.D. | Hasan DEMIROGLU | Celal Bayar University, TR |
| Prof. Dr. | Ugur AVCIBASI | Celal Bayar University, TR |
| Asc. Prof. Dr. | Mehmet ONDER | Celal Bayar University, TR |
| Prof. Dr. | Hasan Huseyin UGURLU | Gazi University, Faculty of Education, Ankara, Turkey |
| Asist. Prof. Dr. | Evren ZIPLAR | Cankiri Karatekin University, TR |
| Assist. Prof. Dr. | Bahriye SIRAV | Gazi University, TR |
| Prof. Dr. | Iskender REYHANCAN | Istanbul Thecnical University, TR |
| Prof. Dr. | Mehmet Ali KORPINAR | Istanbul University, TR |
| Asist. Prof. Dr. | Mehmet TURKMEN | Hacettepe, Dept. of Nuclear Eng., Ankara, Turkey |
| Asist. Prof. Dr. | Tahir CAKIR | Yuzuncu Yıl Universtiy, Van |
| Prof. Dr. | Tunaya KALKAN | Istanbul University, TR |
| Prof. Dr. | Erdal ÇOKÇA | Middle East Technical University TR |
| Asc. Prof. Dr. | Yeliz YUKSELEN AKSOY | Dokuz Eylul University TR |
| Prof. Dr. | Jeffrey C. EVANS | Bucknell University, Pennsylvania, USA |
| Prof. Dr. | Adnan YAZICI | Middle East Technical Univ. (METU) TR |
| Asist. Prof. Dr. | Huseyin BALCI | Adnan Menderes Univ. Aydin, TR |
| Prof. Dr. | Faruk BOZOGLU | Middle East Technical University, (METU) TR |
| Asist. Prof. Dr. | Balati KUERBANJIANG | University of York, York, UK |
| Prof. Dr | Ian WATSON | Systems Power and Energy, Glasgow, UK |

Lycia Press Inc. Office

| Prof. Dr. | Sıddık KESKİN | YYU Faculty of Medicine, Dept. of Medical Statistics. Statistical Editor |
|------------------------------------|---|--|
| Asist. Prof. Dr. | Hakan ERGİN | Istanbul University, Dept. of Foreign Languages. Language Editor |
| Assist. | Elena JALBA | Reading University, London UK. Language Editor |
| Assist | Gonul OZGOK | Lycia Pess Inc. London |
| Typist | Office Lycia Press | office@lycians.com |
| Web | Web Lycia Press | info@lycians.com |
| Assist. Assist Typist Web | Elena JALBA Gonul OZGOK Office Lycia Press Web Lycia Press | Reading University, London UK. Language Editor Lycia Pess Inc. London office@lycians.com info@lycians.com |

- Important
- NSD journal team, is committed to deterring plagiarism, including self-plagiarism. Your manuscripts will be screened for similarity detection with iThenticate, Similarity rate is expected under the %30 except for material and method section.
- For research studies using human or animal subjects, the trial's design, conduct and reporting of results must conform to Good Clinical Practice guidelines (such as the Good Clinical Practice in Food and Drug Administration (FDA)-Regulated Clinical Trials (USA) or the Medical Research Council Guidelines for Good Clinical Practice in Clinical Trials (UK)) and/or to the World Medical Association (WMA) Declaration of Helsinki
- Dear Authors, please load just these three file to manuscript submission system
- Title Page Sample
- Manuscript Sample
- Copyright Transfer and Author Consent Form
- NSD uses vancouver reference style, please prepare articles due to Vancouver reference style rules.
- Manuscript Preparation Rules
- 1.Cover letter
- a- A statement that the manuscript has been read and approved by all the authors.
- b- Approval of all the authors regarding the order in which their names have appeared.
- c- That each author confirms the manuscript represents honest work.
- d- The name, address, and telephone number of the corresponding author who is responsible for communicating with other authors about revisions and final approval.
- e- The letter should give any additional information that may be helpful to the editor, such as the type or format of the article. If the manuscript has been submitted previously to another journal or in another language, it is helpful to include the previous editor's and reviewers' comments with the submitted manuscript, along with the authors' responses to those comments. Submitting previous evaluatory review of another journal accelerates the review process.
- f- For accepted manuscripts, the authors are requested to fill and sign the journal's cover letter to express their consent for its publication.
- g- To reproduce published material, to use illustrations or tables or report information about identifiable people, the author should submit a copy of the permission with the manuscript to the journal.
- 2.Top Ethic Committee Approval
- Inclusion of the approval letter from the relevant Ethics Committee or Institution's Review Board regarding the research protocol and the rights of the subjects (if applicable to the study)
- 3.Top Consent Form
- Attach a copy of the consent form to the letter, if applicable. Consent forms would be evaluated by the Ethics Committee and then signed by the participant.
- 4.Top RCT or NCT Registration
- Emailing the letter denoting registration of RCTs or NCTs in domestic or international databases (The trial's registration number needs to be mentioned, too).
- 5. Manuscripts submitted in English, must be type written, double-spaced, on good quality A4 paper, or paper of similar format. Authors are requested to reserve margins of at least 2.5cm all around the paper. Original drawings of photos, tables and figures should be furnished together with the manuscripts.
- 6. Manuscripts should be kept to a minimum length and should be subdivided into labeled sections (Title page, Abstract, Keywords, Introduction, Materials and Methods, Results, Discussion, Conclusion, Acknowledgement, and References).
- 7. A title page is to be provided and should include the title of the article, authors' names with full first name (with degrees), authors' affiliation, suggested running title and corresponding author. The affiliation should comprise the department, institution (usually university or company), city and state (or nation). The suggested running title should be less than 50 characters (including spaces) and should comprise the article title or an abbreviated version thereof. For office purposes, the title page should include the name and complete mailing address, telephone and fax number, and email of the one author designated to review proofs.
- 8. An abstract no longer than 250 words for reviews and research articles is to be provided as the second page. Abstract should be structured as objective(s) (including purpose setting), materials and methods, results, and conclusion.

Case Report

- A case report is a case study, case report, or other description of a case that should contain 1500 2000 words with a structured abstract of 200 words maximum. Case reports should comprise sections of Introduction, Case Presentation, and Conclusions in Abstract and Introduction, Case Presentation, and Discussion in full text with not more than 2 tables or figures and up to 20 references.
- Brief Report
- Brief Reports should contain 1000 2000 words with a structured abstract of 200 words maximum. Short reports should
 comprise sections of Background, Objectives, Materials & Methods, Results and Discussion with not more than 2 tables or
 figures and up to 20 references.
- Short Communication
- Short Communication, follow the instructions for original articles, except that the total word number of the main text (excluding references, tables and figure legends) is limited to 2000 with no more than 2 figures and/or tables and no more than 15 references. An abstract, not exceeding 150 words, should be presented at the beginning of the article.
- News
- News should contain 1000 2000 words with a structured abstract of 200 words maximum. News should comprise sections of Background, Objectives, Materials & Methods, Results and Discussion with not more than 2 tables or figures and up to 20 references.

Publication Policies

- Manuscripts, or the essence of their content, must be previously unpublished and should not be under simultaneous consideration by another Journal. The authors should also declare if any similar work has been submitted to or published by another Journal. By virtue of the submitted manuscript, the corresponding author acknowledges that all the co-authors have seen and approved the final version of the manuscript. The corresponding author should provide all co-authors with information regarding the manuscript, and obtain their approval before submitting any revisions. Manuscripts are only accepted for publication on the understanding that the authors will permit editorial amendments, though proofs will always be submitted to the corresponding author before being sent finally to press. Prior to the initial submission of a new manuscript, please carefully consider that all authors' names are included as no change to authors' details will be permitted after the acceptance. The decision to accept a contribution rests with the Editorial Board of the NSD.
- Manuscripts will be considered for publication in the form of original articles, Case report, short communications, Letter to editor and review articles. The work should be original or a thorough by an authoritative person in a pertinent field.
- Peer review process
 - All submissions will be reviewed anonymously by at least two independent referees. All manuscripts will be acknowledged upon presenting to the Journal office, provided that all stated requirements are met. Authors are encouraged to suggest names of three expert reviewers, but selection remains a prerogative of the Editor. The whole review process depends on receiving referees comments and revising the manuscripts based on these comments to the author. On receipt of the revised article from the author, and after final approving by referees, the letter of acceptance is issued to the author. Authors have the right to communicate to the editor if they do not wish their manuscript to be reviewed by a particular reviewer because of potential conflicts of interest. No article is rejected unless negative comments are received from at least two reviewers. **NSD employs double blind reviewing process, where both the referee and author remain anonymous throughout the process**.



• Ethical Rules and Rights

• Conflicts of interest

- Conflicts of interest arise when authors, reviewers, or editors have interests that are not fully apparent and that may influence their judgments on what is published. They have been described as those which, when revealed later, would make a reasonable reader feel misled or deceived. (The Committee on Publication Ethics (COPE) states in its Guidelines on Good Publication Practice 2003).
- Authors should disclose, at the time of submission, information on financial conflicts of interest or other interests that may influence the manuscript. Authors should declare sources of funding for the work undertaken.

• The Journal's Policy on Plagiarism

• Any practice of plagiarism will not be tolerated by the journal regarding submitted manuscripts. Non-identifiable quoted segments of articles or close paraphrases from other author/s or even submitting the author's previously published work are known as the act of plagiarism by this journal unless proper use of quotations or paraphrasing with decent citation or referencing are in place. Heavy use of one or a couple of articles is discouraged, even if paraphrased fully. Advertent practice of plagiarism will abort reviewing process or later submission to this journal. All submitted articles will evaluate by *iThenticate* software belonged to cross check for stop any plagiarism and improve publication quality.

• Statement of Human and Animal Rights

- All submitted articles involving human experiments should be performed only in accordance with the ethical standards provided by the responsible committee of the institution and in accordance with the Declaration of Helsinki (as revised in Edinburgh 2000), available at http://www.wma.net/en/30publications/10policies/b3/index.html. Papers describing animal experiments can be accepted for publication only if the experiment conforms the National Institute of Health Guide (National Institute of Health Publications No. 80-23, Revised 1978) for the care and use of Laboratory Animals for experimental procedure. Authors must provide a full description of their anesthetics and surgical procedures. All manuscripts reporting the results of experimental investigations involving human subjects should include a statement confirming the informed consent was obtained from each subject or subject's guardian.
- **Humans:** When reporting experiments on human subjects, authors should indicate whether the procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of 1975, as revised in 2008 (5). If doubt exists whether the research was conducted in accordance with the Helsinki Declaration, the authors must explain the rationale for their approach and demonstrate that the institutional review body explicitly approved the doubtful aspects of the study.
- Animals: When reporting experiments on animals, authors should indicate whether the institutional and national guide for the care and use of laboratory animals was followed.
- All animal or human subjects should be used after approval of the experimental protocol by a local ethics committee.
- Acknowledgements
- Contributors: In acknowledgement section, name people for their contributions or their permission to reproduce their published material, to use their illustrations or provide information about them- try to fully name people who have helped from the conception of the idea to adoption of the hypothesis, to finalization of the study, etc., earnestly.Statement of financial support: Aside from the title page, state any financial or other relationships that might lead to a conflict of interest.
- Copyright
- After acceptance and publication; all ownership rights and Copyrights of the manuscript, passes to international journal of Natural Science and Discovery. Please complete copyright form and send via email to editor. <u>Download NSD Copyright</u> <u>Transfer and Author Consent Form</u>
- •
- This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.
- Copyright 2016: The Author(s); This is an open-access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. All Rights reserved by international journal of Medical Science and Discovery.
- Disposal of material
- Once published, all draft copies of the manuscript, correspondence and artwork will be held at least for 6 months before disposal. Authors and Readers may find original PDF file of article on backup servers such as CLOKKS (https://www.clockss.org/)
- Digital Object Identifier DOI
- Once a manuscript is accepted for publication it will be provided with a registered DOI number following the acceptance decision. Manuscripts accepted for publication by the **NSD** will be published as ahead of print articles prior to the printing date of their scheduled issue. Corresponding author will be provided with a PDF Proof by the publisher once the production process of an accepted manuscript is over.

Instruction for Authors

Article Processing Charge

- NSD is a non-profit Scientific Journal Platform; however, it uses professional services such as Language Editing, DOI, domain and hosting, iThenticate Plagiarism or similarity detection Detection Software. All of these professional services are used for all the article processes and an inevitable cost arises with this.
- Unfortunately, like most open journals, fees of the publication with NSD are charged to Authors. Payment is under the responsibilities of corresponding Author(s). NSD does not charge any fee during the submission period. However, after the peer-review process, a non-refundable charge (100 USD) for each accepted manuscript must be paid by the author(s) via NSD's official PayPal account. An invoice will be sent for each accepted manuscript to corresponding author(s).
- Following with completion of payment procedure, the galey proof and acceptance letter of article will be send to authors for last check
- Preperation of articles in PDF and HTML format is covered by Lycia Press Inc. (press.lycians.com) and Article Processing Charges paids to Lycia Press Inc. (press.lycians.com)
- NSD revenue sources and Sponsorships
- All costs arising from the publications are covered by the Sponsor Companies and Article Processing Charges. Sponsorship request evaluates by the NSD Journal Management Board and the **sponsor company logos** will be included on the back page of printed magazine and in the sponsor section of journal website

| | Article Processing Charge (APC) | Discount % |
|------------------------------------|---------------------------------|------------|
| Regular | 100 USD | |
| for Editorial Board Members | 70 USD | 30% |
| for Affiliated Institution Members | 80 USD | 20% |

• ***APC** not includes Proofreading Services fee. Editor in Chief may direct the corresponding Author to Lycia Press, Language Office for Proofreeading Service<u>lycians.com</u>

References

- Committee on Publication Ethics (COPE). (2011, March 7). Code of Conduct and Best-Practice Guidelines for Journal Editors. Retrieved fromhttp://publicationethics.org/files/Code_of_conduct_for_journal_editors_Mar11.pdf
- World Association of Medical Editors (WAME). Principles of Transparency and Best Practice in Scholarly Publishing. http://www.wame.org/about/principles-of-transparency-and-best-practice

Contents

Original Article

Karyotype Analysis of Tulipa pulchella L. (Liliaceae) (Fenzl ex Regel) Baker
 Yasar Kiran, Gulden Dogan, Zeynep Demirkan
 Natural Science and Discovery, 2016; 2(4): 62-7



Natural Science and Discovery 2016; 2(4):62-7

Original Article

ISSN: 2149-6307

DOI: 10.20863/nsd.63926

Karyotype Analysis of Tulipa pulchella (Liliaceae) (Fenzl ex Regel) Baker

Yasar Kiran^{1*}, Gulden Dogan¹, Zeynep Demirkan¹

Abstract

Objective: In this study the somatic chromosome numbers and detailed morphometric properties of *Tulipa pulchella* (Fenzl ex Regel) Baker was examined.

Material and Methods: The seeds were germinated on moist filter paper in petri dishes at 25°C. Actively growing root tips were pretreated with 0.05 M colchicine for 6 h at room temperature. Then, the root tips were fixed with acetic alcohol (1:3 glacial acetic acid–absolute ethanol) for at least 24 h at 4°C, hydrolyzed in 1 N HCl at 60°C for 1-2 min, then rinsed in tap water for 3–5 min. Finally, they were stained in Feulgen for 1 h and mounted in 45% acetic acid. Digital microphotographs from at least five well-spread metaphase plates were taken using an Olympus BX51 microscope X40 objective and were recorded with an Olympus Camedia C-4000 digital camera.

Results: The chromosome number of *Tulipa pulchella* is determined as 2n = 24 for this taxon. The karyotype consists of 8 submedian region (sm) and 4 subterminal (st) chromosomes. The metaphase chromosome length ranges from 9.35 to 16.48 µm, longest to shortest chromosome ratio was 1.9:4.1, total karyotype length (TKL) was 150.99 um and the karyotype symmetry was type 1A.

Conclusion: The basic chromosome number of *Tulipa pulchella* is x=12. The majority of tulip species and cultivars is diploid (2n=2x=24), however, encounters to triploids (2n=3x=36), tetraploids (2n=4x=48) and even some pentaploids (2n=5x=60) and hexaploid (2n=6x=72). Identifying the chromosome number of this species in this study would provide a base for biosystematic studies.

Keywords: Tulipa pulchella; Chromosome Number; Karyotype Analysis

Introduction

Tulipa L. is a member of Liliaceae, which was formerly treated as a large family, Liliaceae sensu lato. According to the recent phylogenetic studies, Liliaceae includes 15 genera (1-4). The highest diversity of the genus Tulipa is found in the Tien Shan and Pamir Alai mountain ranges in Central Asia. The genus extends from western China, Inner Mongolia and the western Himalaya throughout Central Asia and Iran to the Caucasus, the Balkans, Greece, Cyprus and Turkey, south to the Arabian Peninsula and southeastern Pakistan, and north to the Ukraine and southern Siberia. It extends westwards throughout North Africa to Morocco and the Iberian Peninsula (5). Although the genus is also found throughout the Mediterranean and most other parts of Europe, these regions are not part of the natural distribution of the genus. These tulip species were brought by merchants and travelers from Anatolia and Central Asia to

Europe and escaped from cultivation into the wild where they naturalized (5-11).

Tulipa comprises about 40 or more than 100 species throughout the world, and is widely used as an ornamental plant because of its very attractive flowers. Although European people first introduced Tulipa species with the gift from Kanuni Sultan Süleyman, a sultan of Ottoman Empire, to Emperor Ferdinand I in 1554–1562 today about 65% of the world's cultivated tulips are produced in the Netherlands and the tulip has become a national symbol of this country (12-14). The genus has historically been subdivided into two Tulipa subgenera, (Leiostemones Boiss.) and Eriostemones Boiss., which are clearly distinguished by morphological characteristics (15). Tulipa is represented by 18 taxa in Anatolia and these occur mostly in the Irano-Turanian floristic region of Turkey (16-18).



Received: 27-10-2016 Accepted 30-11-2016 Available Online: 15-12-2016

¹ Firat University, Science Faculty, Dept. of Biology, Elazig, TR

^{*} Corresponding Author: Yasar Kiran E-mail: ykiran@firat.edu.tr

The total number of tulip species is not known exactly, but according to various researchers the genus *Tulipa* is represented by 40–100 species. In the World checklist of selected plant families (19) 503 names have been listed for *Tulipa*, but only 121 taxa have been accepted. Most recently Christenhusz et al. accepted 76 species. Most species are distributed in Central Asia and the Caucasus (65 species), Iran and adjoining regions (36 species) and Turkey (18 species) (4,5).

The genus Tulipa is currently the most important bulbous geophyte in the World and contains around 35 species in the subg. Tulipa with glabrous stamens, and 20 species in the subg. Eriostemones with bossed, usually hairy stamens (5,11,14). Tulipa was revised for the "Flora of Turkey and the east Aegean Islands" by Marais, accepting 14 species and one infaspecific taxon. Since then, three new species have been described from Turkey (17,18,20,21). In Turkey, Eker et al. revised the genus Tulipa. Due to detailed morphologic, geographic and cytotaxonomic studies of the taxa, the genus Tulipa was divided into two subgenera and they represented 17 species, two subspecies and two varieties (in total 19 taxa) (4). Marais reduced T. pulchella (Fenzl ex Regel) Baker to a synonym of T. humilis Herbert, and similarly Van Raamsdonk and De Vries accepted this species as a form of T. humilis (15,18). Christenhusz et al. treated it as a natural variety of T. humilis (5). Although both species show large variation in flower colour, there is consistency of how this varies; mauve in T. humilis and pinkish-purple to magenta colours in T. pulchella are dominant and typical. T. pulchella differs also from T. humilis in having blackish-brown bulb tunics and a bicolored floral blotch. Additionally, these species have an allopatric distribution, at least in Turkey (4). Eker et. al. reported as chromosome numbers of T. pulchella and synonym T. humilis 2n=2x=24 (4).

Material and Methods

Plant material was collected from natural habitats during the fruiting season in Mersin in 2015. Voucher specimen was deposited at the Firat University Herbarium (FUH). Karyological studies were conducted on meristematic cells obtained from the root tips.

The seeds were germinated on moist filter paper in petri dishes at 25°C. Actively growing root tips were pretreated with 0.05 M colchisin for 6 h at room temperature. Then, the root tips were fixed with acetic alcohol (1:3 glacial acetic acid–absolute ethanol) for at least 24 h at 4°C, hydrolyzed in 1 N HCl at 60°C for 1-2 min, then rinsed in tap water for 3–5 min. Finally, they were stained in Feulgen for 1 h and mounted in 45% acetic acid. Digital microphotographs from at least five well-spread metaphase plates were taken using an Olympus BX51 microscope X40 objective (Olympus Optical Co. Ltd., Tokyo, Japan), and were recorded with an Olympus Camedia C-4000 digital

Natural Science and Discovery 2016; 2(4):62-7

camera (Olympus Optical Co. Ltd., Tokyo, Japan). The short arm (S), long arm (L) and total lengths of each chromosome were measured and the relative lengths, arm ratios, and centromeric indices were from images of selected cells. determined Chromosomes were classified according to the nomenclature of Levan et al. (22). The intrachromosomal asymmetry index (A1) and the interchromosomal asymmetry index (A2) followed those of Romero-Zarco (23). The karyotype symmetry nomenclature followed Stebbins (24). Also, relevant literature the online chromosome number databases, Index to Plant Chromosome Numbers (IPCN) (25).



Figure 1: Habit of *T. pulchella* in the wild.

Results

The results of this study showed that the chromosome number of *T. pulchella* is 2n=24. Karyotype analysis of this species to reveal the many values were calculated. The number of somatic chromosome, ploidy level, karyotype formula, chromosome length range, total karyotype length (TKL), Stebbins and asymmetry indexes (A1, A2) are presented in Table 1; relative length, arm ration, centromeric index, type, in Table 2. Metaphase chromosomes in Fig. 2 shown and haploid idiograms of *T. pulchella* in Fig. 3. The basic chromosome number determined as x = 12 for

this taxon. The karyotype consists of 8 submedian region (sm) and 4 subterminal (st) chromosomes. The metaphase chromosome length ranges from 9.35 to 16.48 μ m, longest to shortest chromosome ratio is 1.9:4.1, total karyotype length (TKL) 150.99 μ m and the karyotype symmetry is type 1A.



Figure 2. Somatic metaphases of *T. pulchella*. Scale bar $10 \ \mu m$.

Chromosomes of tulips have been studied since the early twentieth century. Guignard, who first determined the chromosome complement in tulips, found three species to be diploids with 24 pairs of chromosomes (26). Examining varieties of garden tulips, De Mol found a basic number of 12 (27). Also, he discovered one triploid, 'Pink Beauty', with 36 chromosomes. Newton published two papers in which he recorded triploids, tetraploids and one pentaploid (28, 29).Morever, Kroon and Jongerius first discovered a hexaploid origin for T. polychroma Stapf (= T. biflora Stapf) (30). Although many tulips have 24 pairs of chromosomes, some have more than the basic number of 2n=24, including triploids (2n=36), tetraploids (2n=48), pentaploids (2n=60) and exceptionally hexaploids (2n=72) (10,29,31).

Chromosome numbers have been reported for 63 *Tulipa* species (30) and over 600 varieties (32). The basic chromosome number is x=12. The majority of tulip species and cultivars is diploid (2n=2x=24), however, triploids (2n=3x=36), tetraploids (2n=4x=48) and even some pentaploids (2n=5x=60) and hexaploid (2n=6x=72) ocur (30,32-36).

Table 1. Somatic chromosome number, ploidy level, karyotype formula, chromosome length range, total karyotype length (TKL), asymmetry indexes (A1, A2) of Romero Zarco (1986) and symmetry classes (SC) of Stebbins (1971) of *T. Pulchella*.

| Taxon | 2n | Ploidy level | Karyotype formula | Chromosome length range (µm) | TKL (µm) | A1 | A2 | SC |
|------------------|----|-----------------|----------------------|---------------------------------|----------|------|------|----|
| Tulipa pulchella | 24 | 2x | 8sm+4st | 9.35-16.48 | 150.99 | 0.61 | 0.18 | 1A |

| Chromosome Pair No | Total Length µm) | Long Arm(L) µm) | Short Arm(S) µm) | Arm Ration (L/S) | Centromeric Index (CI=S/CL) | Relative Length RL%=(CL/ΣCL)x100 | Chromosome Type |
|-----------------------|------------------------|-----------------------|------------------------|------------------------|-----------------------------------|-------------------------------------|--------------------|
| 1 | 16.48 | 13.26 | 3.22 | 4.11 | 0.19 | 10.92 | st |
| 2 | 15.67 | 10.62 | 5.04 | 2.10 | 0.32 | 10.37 | sm |
| 3 | 15.30 | 12.17 | 3.13 | 3.88 | 0.20 | 10.13 | st |
| 4 | 13.44 | 10.53 | 2.90 | 3.62 | 0.21 | 8.90 | st |
| 5 | 14.03 | 9.26 | 4.76 | 1.94 | 0.33 | 9.29 | sm |
| 6 | 12.62 | 8.99 | 3.63 | 2.47 | 0.28 | 8.36 | sm |
| 7 | 11.12 | 8.67 | 2.45 | 3.53 | 0.22 | 7.37 | st |
| 8 | 11.58 | 8.35 | 3.22 | 2.59 | 0.27 | 7.67 | sm |
| 9 | 10.99 | 8.08 | 2.90 | 2.78 | 0.26 | 7.28 | sm |
| 10 | 10.35 | 7.49 | 2.86 | 2.61 | 0.27 | 6.85 | sm |
| 11 | 9.99 | 6.67 | 3.31 | 2.01 | 0.33 | 6.61 | sm |
| 12 | 9.35 | 6.22 | 3.13 | 1.98 | 0.33 | 6.19 | sm |

 Table 2. Karyo-morphological parameters of T. pulchella.



Figure 3. Haploid idiogram of T. pulchella.

Abedi et al. were examined chromosomal and karyotype parameters and genome size in 22 Iranian populations of nine different *Tulipa* species. Most species were diploid 2n = 2x = 24 (37). 2n=24 chromosomes of this species was reported in the literature (15,38,39). In cytotaxonomic analyses, the basic chromosome number was found to be 2n=24 (x=12) (*T. humilis, T. pulchella*). Polyploidy was observed in some taxa, such as 2n=3x=36 (*T. aleppensis* Boiss. ex Regel, *T. orphanidea* Boiss. ex Heldr), 2n=4x=48 (*T. sylvestris* subsp. *sylvestris* L.) and 2n=5x=60 (*T. clusiana* Red) (4). Therefore, the present count confirmed the earlier reports on 2n chromosomes number.

Conclusion

According to our knowledge, chromosome number and morphology report for *T. pulchella*. Identifying the chromosome number of this species in this study provides a base for biosystematic studies. **Conflict of interest:** The authors declare they have no potential conflicts of interest with respect to the research, authorship, and/or publication of this article, and declare study has ethical permissions if required.

Acknowledgement: Contribution of authors; YK, GD, ZD: Concept and design, YK, GD, ZD: Data Analysis, GD: Writing, GD: Editing

References

- Ronsted N, Law S, Thornton H, Fay MF, Chase MW. Molecular phylogenetic evidence for the monophyly of Fritillaria and Lilium (Liliaceae; Liliales) and the infrageneric classification of Fritillaria. Mol Phylo and Evo. 2005;35:509–527.
- Fay MF, Chase MW, Ronsted N, Devey DS, Pillon Y, Pires JC, et al. Phylogenetics of Liliales: summarized evidence from combined analyses of five plastid and one mitochondrial loci. Aliso. 2006;22: 559–565.

- Angiosperm Phylogeny Group [APG]. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III. Bot J of the Linn Soci. 2009;161:105–121.
- Eker I, Babaç MT, Koyuncu M. Revision of the genus Tulipa L. (Liliaceae) in Turkey. Phytotaxa. 2014;157 (1):001-112.
- Christenhusz MJM, Govaerts R, David JC, Hall T, Borland K, Roberts PS, et al. Tiptoe through the tulips-cultural history, molecular phylogenetics and classification of Tulipa (Liliaceae). Bot J of the Linn Soci. 2013;172:280– 328.
- Hoog MH. On the origin of Tulipa. In: Napier E. and Platt JNO. (eds), Lilies and other Liliaceae. The Royal Horticultural Society: London; 1973. p. 47–64.
- 7. Pavord A. The tulip. Bloomsbury: London; 1999. p. 388.
- Dobbs L. Tulip 70 stunning varieties of the world's favorite flower, St Martin's: New York; 2003. p.104.
- Juodkaite R, Naujalis JR, Navalinskiene M, Samuitiene M. Evaluation of tulip (Tulipa L.) decorative capacities and resistance to tulip breaking potyvirus in the tulip collection of the Botanical Garden of Vilnius University. Biologia. 2005;4:64–70.
- Wilford R. Tulips species and hybrids for the gardener. Timber Press: Portland; 2006. p. 211.
- Tuyl JM Van, Creij MGM Van. Tulipa gesneriana and tulip hybrids. In: Anderson NO, Flower breeding and genetics: issues, challenges, and oppurtunities for the 21st century 23, Springer: Dordrecht; 2006. p. 613–637.
- Ozturk M, Akcicek E, Ozcelik H, Sayar A. Tulip in Turkish Art and Folklore. Proceedings of the second International Scientific Conference: Cario 17–20 March; 1997. p. 367–374.
- 13. Baytop A. Turkiye'de Botanik Tarihi Arastırmaları. TUBITAK Yayınları, Akademik Dizi 3: Ankara; 2004.
- Booy G, Van Raamsdonk LWD. Variation in the enzyme esterase within and between Tulipa species; usefulness for the analysis of genetic relationships at different taxonomical levels. Biochem Syst Ecol. 1998;26:199–224.
- Raamsdonk LWD, Vries T. Biosystematic studies in Tulipa sect. Eriostemones (Liliaceae). Plant Syst and Evol. 1992;179:27–41.
- Guner A, Ozhatay N, Ekim T, Baser KHC. Flora of Turkey and the East Aegean Islands, vol. 11 (Suppl 2), Edinburgh University Press: Edinburgh; 2000.
- Terzioğlu S, Coşkunçelebi K. Tulipa gumusanica (Liliaceae), a new species from Turkey. Annales Botanici Fennici. 2002;39:149–151.
- Marais W. Tulipa. In: Davis PH, Mill RR, Tan K. (eds). Flora of Turkey and the east Aegean Islands, vol. 8, Edinburgh University Press: Edinburgh; 1984. p. 245–263.

- Govaerts R. Tulipa. In: Govaerts R. (ed). World checklist of selected plant families: RBG Kew: UK; 2013.
- Persson K. Two new bulbous species from the central Taurus Mountains of Turkey. New Plantsman. 2000;7:200– 208.
- Özhatay N, Koçak B. Tulipa karamanica. In: Güner A, Özhatay N, Ekim T, Başer KHC. (eds). Flora of Turkey and the east Aegean Islands, vol. 11(Suppl. II). Edinburgh University Press: Edinburgh; 2000. p. 246.
- 22. Levan A, Fredga K, Sanberg AA. Nomenclature for centromeric position on chromosomes. Hereditas. 1964;52(2):201–220.
- 23. Romero-Zarco C. A new method for estimating karyotype asymmetry. Taxon. 1986;35: 526–530.
- 24. Stebbins GL. Chromosome Evolution in Higher Plants. Edward Arnold: London;1971.
- 25. http://www.tropicos.org/Project/IPCN
- Guignard L. L'appareil sexuel et la double fécondation dans les tulipes. Ann des Sci Nat Botanique (ser. viii). 1900;11:365–387.
- Mol WE de. Het celkundig-erfelijk onderzoek in dienst gesteld van de veredeling der hyacinthen, narcissen en tulpen. Genetica. 1925;7:111–118.
- Newton WCF. Chromosome studies in Tulipa and some related genera. J of the Linn Soci Bot. 1927;47:339–354.
- Newton WCF, Darlington CD. Meiosis in polyploids. I. Triploid and pentaploid tulips. J of Genetics. 1929;21: 1– 15.
- Kroon GH, Jongerius MC. Chromosome numbers of Tulipa species and the occurrence of hexaploidy. Euphytica. 1986;35:73–76.
- Hall AD. The genus Tulipa. Royal Horticultural Society: London; 1940. p.171.
- Zeilinga AE, Schouten HP. Polyploidy in garden tulips. I. Survey of Tulipa varieties for polyploids. Euphytica. 1968a;17:252-264.
- Zeilinga AE, Schouten HP. Polyploidy in garden tulips. II. The production of tetraploids. Euphytica. 1968b;17:303-310.
- Holitscher O. Pruhonicky sortiment tulipanu. Acta Pruhoniciana. 1968;18:1-215.
- Kroon GH. Choromosome number of garden tulips. Acta Bot Neerlandica. 1975;24: 489-490.
- Van Scheepen J. Classified list and international register of tulip names. Royal General Bulbgrowers Association KAVB, Hillegom: The Netherlands; 1996.
- Abedi R, Babaei A, Karimzadeh G. Karyological and flow cytometric studies of Tulipa (Liliaceae) species from Iran. Plant Syst Evol. 2015; 301: 1473–1484.

- Southern DI. Species relationship in the genus Tulipa. Chromosoma. 1967;23:80-94.
- Blakey DH, Vosa CG. Heterochromatin and chromosome varition in cultivated species of Tulipa, subg. Eriostemones (Liliaceae). Plant Syst and Evol. 1981;139:47-55.

Copyright © 2016 The Author(s); This is an open-access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. All Rights reserved by international journal of Natural Science and Discovery



NATURAL SCIENCE & DISCOVERY

ISSN: 2149-6307 Lycia Press LONDON UK



International Journal of Natural Science and Discovery Open Access Scientific Journal 2016, Vol.2, No.4



www.natscidiscovery.com